

09/972,177

=> d his

(FILE 'HOME' ENTERED AT 12:32:05 ON 04 NOV 2003)

FILE 'REGISTRY' ENTERED AT 12:32:15 ON 04 NOV 2003

L1 STRUCTURE UPLOADED

L2 QUE L1

L3 7 S L2

L4 1344 S L2 SSS FUL

FILE 'CAPLUS' ENTERED AT 12:34:24 ON 04 NOV 2003

L5 133 S L4

FILE 'REGISTRY' ENTERED AT 12:36:05 ON 04 NOV 2003

L6 STRUCTURE UPLOADED

L7 QUE L6

L8 STRUCTURE UPLOADED

L9 QUE L8

L10 254 S L7 SUB=L4 FUL

L11 113 S L9 SUB=L4 FUL

L12 347 S L10 OR L11

FILE 'CAPLUS' ENTERED AT 12:37:25 ON 04 NOV 2003

L13 64 S L12

FILE 'REGISTRY' ENTERED AT 12:37:41 ON 04 NOV 2003

L14 692 S L4 AND NRS>2

L15 200 S L12 AND NRS>2

L16 147 S L12 NOT L15



FILE 'CAPLUS' ENTERED AT 12:42:43 ON 04 NOV 2003

L17 39 S L15

=> d 12

L2 HAS NO ANSWERS

L1 STR



Structure attributes must be viewed using STN Express query preparation.


L2 QUE ABB=ON PLU=ON L1

=> d 17

L7 HAS NO ANSWERS


L6 STR

09/972,177



Structure attributes must be viewed using STN Express query preparation.  
L7 QUE ABB=ON PLU=ON L6

=> d 19  
L9 HAS NO ANSWERS  
L8 STR




Structure attributes must be viewed using STN Express query preparation.  
L9 QUE ABB=ON PLU=ON L8

=> d ibib abs hitstr 1-39 117

09/972,177

L17 ANSWER 1 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 2002:594840 CAPLUS  
 DOCUMENT NUMBER: 137:154858  
 TITLE: Preparation of arylsulfonamidopiperidones as inhibitors of Factor Xa.  
 INVENTOR(S): Stein, Philip P.; O'Connor, Stephen P.; Lawrence, R. Michael; Shi, Yan  
 PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA  
 SOURCE: PCT Int. Appl., 246 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002060894	A2	20020808	WO 2002-US2542	20020128
WO 2002060894	A3	20021219		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
US 6555542	B1	20030429	US 2002-59621	20020129
PRIORITY APPLN. INFO:			US 2001-264964P	P 20010130
OTHER SOURCE(S):		MARPAT 137:154858		
GI				



AB Title compds. [I; X = (substituted) (CH2)<sub>m</sub>; m = 1-3; R1 = (substituted) alkyl, alkenyl, alkynyl, aryl, heteroaryl, etc.; R2, R3 = H, (substituted) alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, etc.; R4, R5, R6, R7, R8 = H, OH, (substituted) alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, alkoxy, etc.; R9, R10 = H, (substituted) alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl, etc.; R11, R12 = (substituted) (CH2)<sub>n</sub>; n = 1-4; R13-R16 = (substituted) cycloalkyl, etc. prep. as cardiovascular agents (no data). 974 I, including (II), were prep.

L17 ANSWER 2 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 2002:314941 CAPLUS  
 DOCUMENT NUMBER: 136:340701  
 TITLE: Preparation of 3,8-diazabicyclo[3.2.1]octanes for treating cardiac arrhythmias  
 INVENTOR(S): Bjoersne, Magnus; Hoffmann, Kurt-Juergen; Ponten, Fritiof; Strandlund, Gert; Svensson, Peder; Wlstermann, Michael  
 PATENT ASSIGNEE(S): AstraZeneca AB, Sweden  
 SOURCE: PCT Int. Appl., 135 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002032902	A1	20020425	WO 2001-SE2294	20011018
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, BG, BR, BU, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2001096173	A5	20020429	AU 2001-96173	20011018
EP 1328526	A1	20030723	EP 2001-977023	20011018
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
PRIORITY APPLN. INFO:			SE 2000-3795	A 20001020
OTHER SOURCE(S):		MARPAT 136:340701	WO 2001-SE2294	W 20011018
GI				



AB The title compds. [I; one of R1 and R2 = R1a and the other = ACR13R14R15 (wherein R1a = alkyl optionally substituted and/or terminated by one or more groups selected from halo, CN, NO2, etc.; R13 = H, halo, alkyl, etc.; R13R14 = O; or R14 = H, alkyl; R15 = (un)substituted aryl, heteroaryl; A = alkylene, etc.; B = a bond, alkylene, etc.); R3-R10 = H, alkyl], useful in the prophylaxis and in the treatment of arrhythmias, in particular atrial and ventricular arrhythmias, were prep. Thus, reacting tert-Bu

L17 ANSWER 1 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (prepn. of arylsulfonamidopiperidones as inhibitors of Factor Xa)  
 RN 445273-60-9 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 3-[(3S)-3-[(6-bromo-2-naphthalenyl)sulfonyl]amino]-2-oxo-1-piperidinyl]acetyl]-8-(phenylmethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



L17 ANSWER 2 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
 3-(4-cyanophenoxy)-1-[(3,8-diazabicyclo[3.2.1]oct-8-ylmethyl)propyl]carbamate (prepn. given) with Bu isocyanate in the presence of Et3N in MeCN followed by treatment with HCl/EtOAc afforded I [R1 = CONH2; R2 = CH2CHNH2CH2CH2O-p-C6H4CN; R3-R10 = H] in quant. yield. The exemplified compds. I showed pIC50 values of at least 5.5 for K channel blockade.  
 IT 415975-69-8P 415975-70-1P 415976-92-0P  
 415977-56-9P 415977-64-9P 415977-72-9P  
 415977-80-9P 415977-86-5P 415977-95-6P  
 415978-02-8P  
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (prepn. of 3,8-diazabicyclo[3.2.1]octanes for treating cardiac arrhythmias)


RN 415975-69-8 CAPLUS  
 CN Benzotrifluoride, 4-[(2S)-2-amino-3-[(3-phenylmethyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]propoxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 415975-70-1 CAPLUS  
 CN Carbamic acid, [(1S)-1-[(4-cyanophenoxy)methyl]-2-[3-(phenylmethyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]ethyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 415976-92-0 CAPLUS  
 CN Benzonitrile, 4-[1-(3,4-dimethoxyphenoxy)-4-[(3-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]ethyl)]- (9CI) (CA INDEX NAME)


L17 ANSWER 2 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




RN 415977-56-9 CAPLUS  
 CN Benzonitrile, 4-[[3-[[3-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-hydroxypropoxy]-9CI] (CA INDEX NAME)



RN 415977-64-9 CAPLUS  
 CN Benzonitrile, 4-[[3-[[3-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]propyl]amino]-9CI] (CA INDEX NAME)



RN 415977-72-9 CAPLUS  
 CN Benzonitrile, 4-[[3-[[3-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]ethoxy]-9CI] (CA INDEX NAME)




RN 415977-80-9 CAPLUS  
 CN Benzonitrile, 4-[[3-[[3-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]propyl]sulfonyl]-9CI] (CA INDEX NAME)


L17 ANSWER 2 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)

IT 17740-41-9P 415979-11-2P 415979-13-4P  
 415979-15-6P 415979-34-9P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (prepn. of 3,8-diazabicyclo[3.2.1]octanes for treating cardiac arrhythmias)


RN 17740-41-9 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-2,4-dione, 3,8-bis(phenylmethyl)-, monohydrochloride (9CI) (CA INDEX NAME)



RN 415979-11-2 CAPLUS  
 CN Benzonitrile, 4-[[2-hydroxy-3-[(3-phenylmethyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]propoxy]-9CI] (CA INDEX NAME)




RN 415979-13-4 CAPLUS  
 CN Benzonitrile, 4-[[3-[[3-[(phenylmethyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]propyl]amino]-9CI] (CA INDEX NAME)




RN 415979-15-6 CAPLUS  
 CN Benzonitrile, 4-[[3-[[3-[(3,4-dimethoxyphenoxy)-4-[(3-phenylmethyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]butyl]-9CI] (CA INDEX NAME)

L17 ANSWER 2 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




RN 415977-86-5 CAPLUS  
 CN 1,3-Benzenedicarbonitrile, 4-[[2-[[3-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]ethoxy]-9CI] (CA INDEX NAME)




RN 415977-95-6 CAPLUS  
 CN Carbamic acid, [(1S)-2-[(4-cyanophenoxy)-1-[[3-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]methyl]ethyl]-, methyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.




RN 415978-02-8 CAPLUS  
 CN Benzonitrile, 4-[[3-amino-4-[[3-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]butoxy]-9CI] (CA INDEX NAME)



L17 ANSWER 2 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



Absolute stereochemistry.




REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

09/972,177

L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 2002:314940 CAPLUS  
 DOCUMENT NUMBER: 136:340711  
 TITLE: Bridged piperazine derivatives, specifically 3,8-diazabicyclo[3.2.1]octane, 8-azabicyclo[3.2.1]octane, 2,5-diazabicyclo[2.2.2]octane, and 3,9-diazabicyclo[3.3.1]nonane derivatives, useful as inhibitors of chemokines binding to CCR1 receptors, for treating inflammation and other immune disorders.  
 INVENTOR(S): Blumberg, Laura Cook; Brown, Matthew Frank; Glaude, Ronald Paul; Poss, Christopher Stanley  
 PATENT ASSIGNEE(S): Pfizer Products Inc., USA  
 SOURCE: PCT Int. Appl., 89 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002032901	A2	20020425	WO 2001-1B1844	20011004
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TH, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
AU 2001092160	A5	20020429	AU 2001-92160	20011004
EP 1326867	A2	20030716	EP 2001-972389	20011004
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
US 200219961	A1	20020829	US 2001-972177	20011005
NO 2003001572	A	20030610	NO 2003-1572	20030408
PRIORITY APPL. INFO.:			US 2000-241804P	P 20001019
			WO 2001-1B1844	W 20011004
OTHER SOURCE(S):		MARPAT 136:340711		
GI				

L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



AB Compds. I and their pharmaceutically acceptable salts, useful for treatment of inflammation and other immune disorders, are disclosed [wherein: n = 1-5; m = 1-5; q = 0-1; a, b, c = (CH2)0-4 (independently); a, b, and c cannot all be null; if a and/or c is not null, then b must be null; W = CH or N; X = CO, C(S), or CH2; Z = O, (un)substituted NH or (un)substituted CH2; R = certain (un)substituted (hetero)aryl or (hetero)cycloalkyl; R1 = (independently) H, OH, SO3H, halo, alkyl, SH, CF3, wide variety of other substituents]. The compds. are useful for treatment of a wide variety of diseases and disorders, which are cited specifically in claims. Approx. 100 specific examples of I are given, many with synthetic details. For example, 3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]octan-2-one (prepn. given) underwent a sequence of: (1) reduct. of the amide carbonyl using LiAlH4 (94%); (2) 8-N-acylation with chloroacetyl chloride (69%); and (3) etherification with 2-nitro-4-trifluoromethylphenol (58%), to give title compd. II. In a bioassay for the ability to inhibit chemotaxis of various cells (THP-1 cells, primary human monocytes, or primary lymphocytes) in vitro, all example compds. had IC50 values of less than 10 μM.

IT 417726-56-EP 417726-79-SP, 2-(2-Amino-5-chlorophenoxymethyl)-1-[3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]ethanone 417726-83-1P, 2-(3-Amino-5-chloropyridin-2-ylomethyl)-1-[3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]ethanone


RL: PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)

(drug candidate; prepn. of bridged piperazine derivs. as inhibitors of chemokines binding to CCR1 receptors)


RN 417726-56-8 CAPLUS

CN Benzamide, N-(2-aminoethyl)-4-chloro-2-[2-{3-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]- (9CI) (CA INDEX NAME)


L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



RN 417726-79-5 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8-[(2-amino-5-chlorophenoxymethyl)-3-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)



RN 417726-83-1 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8-[(3-amino-5-chloro-2-pyridinyl)oxy]acetyl-3-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)




IT 417726-39-7P, 1-[3-(4-Fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-(2-nitro-4-trifluoromethylphenoxymethyl)ethanone 417726-40-OP, 4-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]benzamide 417726-41-1P, 1-[3-(4-Fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-(2-ethoxycarbonyl-4-chlorophenoxymethyl)ethanone 417726-42-2P, 1-[3-(4-Fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-(2-acetyl-5-chlorophenoxymethyl)ethanone 417726-43-3P, 1-[3-(4-Fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-(2-sulfamoyl-5-chlorophenoxymethyl)ethanone 417726-44-4P, 1-[3-(4-Fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-(2-nitro-5-trifluoromethylphenoxymethyl)ethanone 417726-45-5P, 1-[3-(4-Fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-[2-(ethoxycarbonyl)methyl]-5-chlorophenoxymethyl)ethanone 417726-46-6P, 5-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]benzenesulfonamide 417726-47-7P, 4-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]benzenesulfonamide 417726-48-8P, 4-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]benzamide 417726-49-9P,


L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)

5-Methoxy-2-[2-{8-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-3-yl]-2-oxoethoxy]benzamide 417726-50-2P, 4-Chloro-2-[2-{8-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-3-yl]-2-oxoethoxy]-1-nitrobenzene 417726-51-3P, 2-(5-Chloroquinolin-8-ylomethyl)-1-[3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]ethanone 417726-52-4P, 1-[3-(4-Fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-(6-methyl-2-nitropyridin-3-ylomethyl)ethanone 417726-53-5P, 2-(5-Chloro-3-nitropyridin-2-ylomethyl)-1-[3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]ethanone 417726-54-6P, 2-(5-Chloro-3-carbamoylpyridin-2-ylomethyl)-1-[3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]ethanone 417726-55-7P, 2-(5-Chloro-3-methoxycarbonylpyridin-2-ylomethyl)-1-[3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]ethanone 417726-56-8P, 4-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]benzoic acid 417726-58-OP, 4-Methyl-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]benzoic acid 417726-59-1P, 4-Methoxy-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]benzoic acid 417726-60-4P, 4-Iodo-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]benzoic acid 417726-61-5P, 4-Bromo-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]benzoic acid 417726-62-6P, 4-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]benzenesulfonamide 417726-63-7P, 5-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]benzoic acid 417726-64-8P, 5-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]nicotinic acid 417726-65-9P, 3-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]naphthalene-2-carboxylic acid 417726-66-OP, 4-Chloro-1-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]naphthalene-2-carboxylic acid 417726-67-1P, 5-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]-N-(1H-tetrazol-5-yl)benzamide 417726-68-2P, 4-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]-N-(carboxymethyl)benzamide 417726-69-3P, 4-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]-N-(methanesulfonyl)benzamide 417726-70-6P, 4-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]-N-(1H-tetrazol-5-yl)benzamide 417726-71-7P, 4-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]-N-(carbamoylmethyl)benzamide 417726-72-8P, N-Carbamoylmethyl-5-chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]nicotinamide 417726-73-9P, 4-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]-N-(2-ureidomethyl)benzamide 417726-74-OP, 1-[5-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]urea 417726-75-1P, N-[4-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]urea 417726-76-2P, 1-[4-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]phenyl]-3-(2-carboxymethyl)urea 417726-77-3P, 5-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]urea 417726-78-4P, 2-[3-(5-Chloro-2-[2-{3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]pyridin-3-yl)ureido]acetamide 417726-80-8P, 2-(2-Amino-4-trifluoromethylphenoxymethyl)-1-[3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]ethanone 417726-81-9P, 2-(2-Amino-4-chlorophenoxymethyl)-1-[3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]ethanone 417726-82-OP, 2-(2-Amino-4-chlorophenoxymethyl)-1-[8-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-3-yl]ethanone 417726-84-2P, 2-(2-Amino-6-methylpyridin-3-ylomethyl)-1-[3-(4-fluorobenzyl)-3,8-


L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
 diazabicyclo[3.2.1]oct-8-yl]ethanone 417726-85-3P,  
 2-Amino-N-[5-chloro-2-[2-[3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]pyridin-3-yl]acetamide 417726-86-4P,  
 N-[5-Chloro-2-[2-[8-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-3-yl]-2-oxoethoxy]phenyl]-3-hydroxy-3-methylbutyramide 417726-87-5P,  
 N-[4-Chloro-2-[2-[3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]phenyl]methanesulfonamide 417726-88-6P,  
 N-[5-(Trifluoromethyl)-2-[2-[3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]phenyl]methanesulfonamide 417726-89-7P,  
 N-[5-Chloro-2-[2-[3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]phenyl]methanesulfonamide 417726-90-0P,  
 N-[2-[[[4-Chloro-2-[2-[3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]phenyl]carbonyl]amino]ethyl]methanesulfonamide 417726-91-1P,  
 N-[5-Chloro-2-[2-[8-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-3-yl]-2-oxoethoxy]phenyl]methanesulfonamide 417726-92-2P,  
 N-[5-Chloro-2-[2-[3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]pyridin-3-yl]methanesulfonamide 417726-93-3P,  
 N-[6-Methyl-3-[2-[3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]pyridin-2-yl]methanesulfonamide 417726-94-4P,  
 5-Chloro-2-[2-[3-(4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]benzamide  
 RL: PAC (Pharmacological activity); SPN (Synthetic Preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (drug candidate; prepn. of bridged piperazine derivs. as inhibitors of chemokines binding to CCR1 receptors)  
 RN 417726-39-7 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 3-[(4-fluorophenyl)methyl]-8-[[2-nitro-4-(trifluoromethyl)phenoxy]acetyl]- (9CI) (CA INDEX NAME)




RN 417726-40-0 CAPLUS  
 CN Benzamide, 4-chloro-2-[2-[3-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]- (9CI) (CA INDEX NAME)




L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



RN 417726-45-5 CAPLUS  
 CN Benzenesulfonamide, 4-chloro-2-[2-[3-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]-, ethyl ester (9CI) (CA INDEX NAME)




RN 417726-46-6 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 3-[[2-(aminosulfonyl)-4-chlorophenoxy]acetyl]-8-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)




RN 417726-47-7 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 3-[[2-(aminosulfonyl)-5-chlorophenoxy]acetyl]-8-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)


L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
 RN 417726-41-1 CAPLUS  
 CN Benzoic acid, 5-chloro-2-[2-[3-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]-, ethyl ester (9CI) (CA INDEX NAME)



RN 417726-42-2 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8-[[2-(2-acetyl-5-chlorophenoxy)acetyl]-3-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)




RN 417726-43-3 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8-[[2-(aminosulfonyl)-5-chlorophenoxy]acetyl]-3-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)




RN 417726-44-4 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 3-[(4-fluorophenyl)methyl]-8-[[2-nitro-5-(trifluoromethyl)phenoxy]acetyl]- (9CI) (CA INDEX NAME)


L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




RN 417726-48-8 CAPLUS  
 CN Benzamide, 4-chloro-2-[2-[8-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-3-yl]-2-oxoethoxy]- (9CI) (CA INDEX NAME)



RN 417726-49-9 CAPLUS  
 CN Benzamide, 2-[2-[8-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-3-yl]-2-oxoethoxy]-5-methoxy- (9CI) (CA INDEX NAME)




RN 417726-50-2 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 3-[[5-chloro-2-nitrophenoxy]acetyl]-8-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)



L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)


RN 417726-51-3 CAPLUS

CN 3,8-Diazabicyclo[3.2.1]octane, 8-[[5-chloro-8-quinolinyl]oxy]acetyl]-3-[[4-fluorophenyl]methyl]- (9CI) (CA INDEX NAME)




RN 417726-52-4 CAPLUS

CN 3,8-Diazabicyclo[3.2.1]octane, 3-[[4-fluorophenyl]methyl]-8-[[6-methyl-2-nitro-3-pyridinyl]oxy]acetyl]- (9CI) (CA INDEX NAME)



RN 417726-53-5 CAPLUS


CN 3,8-Diazabicyclo[3.2.1]octane, 8-[[5-chloro-3-nitro-2-pyridinyl]oxy]acetyl]-3-[[4-fluorophenyl]methyl]- (9CI) (CA INDEX NAME)



RN 417726-54-6 CAPLUS


CN 3-Pyridinecarboxamide, 5-chloro-2-[2-[3-[[4-fluorophenyl]methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]-4-methyl- (9CI) (CA INDEX NAME)

L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




RN 417726-59-1 CAPLUS

CN Benzoic acid, 2-[2-[3-[[4-fluorophenyl]methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]-4-methoxy- (9CI) (CA INDEX NAME)




RN 417726-60-4 CAPLUS

CN Benzoic acid, 2-[2-[3-[[4-fluorophenyl]methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]-4-iodo- (9CI) (CA INDEX NAME)



RN 417726-61-5 CAPLUS


CN Benzoic acid, 4-bromo-2-[2-[3-[[4-fluorophenyl]methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]- (9CI) (CA INDEX NAME)



RN 417726-62-6 CAPLUS


CN Benzeneacetic acid, 4-chloro-2-[2-[3-[[4-fluorophenyl]methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]- (9CI) (CA INDEX NAME)

L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




RN 417726-55-7 CAPLUS

CN 3-Pyridinecarboxylic acid, 5-chloro-2-[2-[3-[[4-fluorophenyl]methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]-, methyl ester (9CI) (CA INDEX NAME)



RN 417726-57-9 CAPLUS


CN Benzoic acid, 4-chloro-2-[2-[3-[[4-fluorophenyl]methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]- (9CI) (CA INDEX NAME)



RN 417726-58-0 CAPLUS


CN Benzoic acid, 2-[2-[3-[[4-fluorophenyl]methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]-4-methyl- (9CI) (CA INDEX NAME)

L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




RN 417726-63-7 CAPLUS

CN Benzoic acid, 5-chloro-2-[2-[3-[[4-fluorophenyl]methyl]-3,8-diazabicyclo[3.2.1]oct-3-yl]-2-oxoethoxy]- (9CI) (CA INDEX NAME)




RN 417726-64-8 CAPLUS

CN 3-Pyridinecarboxylic acid, 5-chloro-2-[2-[3-[[4-fluorophenyl]methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]- (9CI) (CA INDEX NAME)



RN 417726-65-9 CAPLUS


CN 2-Naphthalenecarboxylic acid, 3-[2-[3-[[4-fluorophenyl]methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]- (9CI) (CA INDEX NAME)




RN 417726-66-0 CAPLUS

CN 2-Naphthalenecarboxylic acid, 4-chloro-1-[2-[3-[[4-fluorophenyl]methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]- (9CI) (CA INDEX NAME)


L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



RN 417726-67-1 CAPLUS  
CN Benzamide, 5-chloro-2-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl-2-oxoethoxy-N-1H-tetrazol-5-yl- (9CI) (CA INDEX NAME)




RN 417726-68-2 CAPLUS  
CN Glycine, N-[4-chloro-2-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxybenzoyl- (9CI) (CA INDEX NAME)




RN 417726-69-3 CAPLUS  
CN Benzamide, 4-chloro-2-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl-2-oxoethoxy-N-(methylsulfonyl)- (9CI) (CA INDEX NAME)


L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



RN 417726-73-9 CAPLUS  
CN Benzamide, N-[[[5-chloro-2-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]phenyl]amino]carbonyl- (9CI) (CA INDEX NAME)




RN 417726-74-0 CAPLUS  
CN Acetamide, 2-[[[5-chloro-2-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]phenyl]amino]carbonyl]amino]- (9CI) (CA INDEX NAME)




RN 417726-75-1 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 8-[[2-[(aminocarbonyl)amino]-5-chlorophenoxy]acetyl]-3-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)


L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



RN 417726-70-6 CAPLUS  
CN Benzamide, 4-chloro-2-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl-2-oxoethoxy-N-1H-tetrazol-5-yl- (9CI) (CA INDEX NAME)




RN 417726-71-7 CAPLUS  
CN Benzamide, N-(2-amino-2-oxoethyl)-4-chloro-2-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl-2-oxoethoxy- (9CI) (CA INDEX NAME)




RN 417726-72-8 CAPLUS  
CN 3-Pyridinecarboxamide, N-(2-amino-2-oxoethyl)-5-chloro-2-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl-2-oxoethoxy- (9CI) (CA INDEX NAME)


L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



RN 417726-76-2 CAPLUS  
CN .beta.-Alanine, N-[[[4-chloro-2-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]phenyl]amino]carbonyl]- (9CI) (CA INDEX NAME)




RN 417726-77-3 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 3-[[2-[(aminocarbonyl)amino]-4-chlorophenoxy]acetyl]-8-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)




RN 417726-78-4 CAPLUS  
CN Acetamide, 2-[[[5-chloro-2-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]-3-pyridinyl]amino]carbonyl]amino]- (9CI) (CA INDEX NAME)




L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




RN 417726-80-8 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 8-[[2-amino-4-(trifluoromethyl)phenoxy]acetyl]-3-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)



RN 417726-81-9 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 8-[(2-amino-4-chlorophenoxy)acetyl]-3-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)




RN 417726-82-0 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 3-[(2-amino-4-chlorophenoxy)acetyl]-8-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)




RN 417726-84-2 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 8-[[2-amino-6-methyl-3-pyridinyl]oxy]acetyl]-3-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)


L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



RN 417726-88-6 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 3-[(4-fluorophenyl)methyl]-8-[[2-[(methylsulfonyl)amino]-4-(trifluoromethyl)phenoxy]acetyl]- (9CI) (CA INDEX NAME)




RN 417726-89-7 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 8-[[4-chloro-2-[(methylsulfonyl)amino]phenoxy]acetyl]-3-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)




RN 417726-90-0 CAPLUS  
CN Benamide, 4-chloro-2-[2-[3-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]-N-[2-[(methylsulfonyl)amino]ethyl]- (9CI) (CA INDEX NAME)


L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



RN 417726-85-3 CAPLUS  
CN Acetamide, 2-amino-N-[5-chloro-2-[2-[3-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]-3-pyridinyl]- (9CI) (CA INDEX NAME)




RN 417726-86-4 CAPLUS  
CN Butanamide, N-[5-chloro-2-[2-[8-[(4-fluorophenyl)methyl]-3,8-diazabicyclo[3.2.1]oct-3-yl]-2-oxoethoxy]phenyl]-3-hydroxy-3-methyl- (9CI) (CA INDEX NAME)




RN 417726-87-5 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 8-[[5-chloro-2-[(methylsulfonyl)amino]phenoxy]acetyl]-3-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)


L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



RN 417726-91-1 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 3-[(4-chloro-2-[(methylsulfonyl)amino]phenoxy]acetyl]-8-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)




RN 417726-92-2 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 8-[[5-chloro-3-[(methylsulfonyl)amino]-2-pyridinyl]oxy]acetyl]-3-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)




RN 417726-93-3 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 3-[(4-fluorophenyl)methyl]-8-[[6-methyl-2-[(methylsulfonyl)amino]-3-pyridinyl]oxy]acetyl]- (9CI) (CA INDEX NAME)


L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS ON STN (Continued)



RN 417728-09-7 CAPLUS  
CN Benzamide, 5-chloro-2-[2-((4-fluorophenyl)methyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy- (9CI) (CA INDEX NAME)



IT 417727-51-6P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(intermediate; prepn. of bridged piperazine derivs. as inhibitors of chemokines binding to CCR1 receptors)  
RN 417727-51-6 CAPLUS  
CN Carbanic acid, [2-[[5-chloro-2-[2-((4-fluorophenyl)methyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy]-3-pyridinyl]amino]-2-oxoethyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)




IT 417727-48-1, 4-Chloro-2-[2-((4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl)-2-oxoethoxy]benzoic acid methyl ester  
417727-49-2, 5-Chloro-2-[2-((4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl)-2-oxoethoxy]benzoic acid 417727-50-5, 2-(5-Chloro-2-nitrophenoxy)-1-[3-((4-fluorobenzyl)-3,8-diazabicyclo[3.2.1]oct-8-yl)]ethanone

L17 ANSWER 4 OF 39 CAPLUS COPYRIGHT 2003 ACS ON STN

ACCESSION NUMBER: 2002:104660 CAPLUS  
DOCUMENT NUMBER: 136:151174  
TITLE: Preparation of 3-[(arylazabicycloalkyl)alkyl]quinazoline-2,4-diones as serotonin reuptake inhibitors and 5-HT2A receptor antagonists  
INVENTOR(S): Butler, Todd William; Filiri, Anton Franz Josef; Gallaschun, Randall James  
PATENT ASSIGNEE(S): Pfizer Products Inc., USA  
SOURCE: Eur. Pat. Appl., 68 pp.  
DOCUMENT TYPE: CODEN: EPXXDW  
LANGUAGE: Patent  
FAMILY ACC. NUM. COUNT: 1 English  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1178048	A1	20020206	EP 2001-306629	20010802
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
US 2002052355	A1	20020502	US 2001-920500	20010801
US 6552015	B2	20030422		
BR 2001003210	A	20020326	BR 2001-3210	20010803
JP 200214769	A2	20020416	JP 2001-236982	20010803
PRIORITY APPL. INFO.:		US 2000-222707P	P	20000803
OTHER SOURCE(S):		MARPAT 136:151174		
GI				




AB R(CH2)nZr1 [1: e.g., (un)substituted 2,4-dioxoquinazolin-3-yl]; R1 = e.g., (un)substituted Ph; Z = azabicycloalkylene; n = 3 or 4) were prepd. Thus, 3,2-cl(H2N)C6H3CO2H underwent cyclocondensation/cyclization with Cl(CH2)3NCO to give 8-chloro-3,4-dihydro-2H-1-oxa-4a,9-diazanthracene-10-one which underwent aminative ring opening with 3-(4-chlorophenyl)-3,8-diazabicyclo[3.2.1]octane to give title compd. II. Data for biol. activity of I were given.  
IT 395059-29-7P 395059-55-9P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(prepn. of 3-[(arylazabicycloalkyl)alkyl]quinazoline-2,4-diones as serotonin reuptake inhibitors and 5-HT2A receptor antagonists)  
RN 395059-29-7 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 3-(4-fluorophenyl)-8-(phenylmethyl)- (9CI) (CA INDEX NAME)


L17 ANSWER 3 OF 39 CAPLUS COPYRIGHT 2003 ACS ON STN (Continued)

RL: RCT (Reactant); RACT (Reactant or reagent)  
(precursor; prepn. of bridged piperazine derivs. as inhibitors of chemokines binding to CCR1 receptors)


RN 417727-48-1 CAPLUS  
CN Benzoic acid, 4-chloro-2-[2-((4-fluorophenyl)methyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy-, methyl ester (9CI) (CA INDEX NAME)




RN 417727-49-2 CAPLUS  
CN Benzoic acid, 5-chloro-2-[2-((4-fluorophenyl)methyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-2-oxoethoxy- (9CI) (CA INDEX NAME)




RN 417727-50-5 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 8-[(5-chloro-2-nitrophenoxy)acetyl]-3-[(4-fluorophenyl)methyl]- (9CI) (CA INDEX NAME)



L17 ANSWER 4 OF 39 CAPLUS COPYRIGHT 2003 ACS ON STN (Continued)




RN 395059-55-9 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 8-(4-chlorophenyl)-3-(phenylmethyl)- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 13 THERE ARE 13 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 5 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 2002:104659 CAPLUS  
 DOCUMENT NUMBER: 136:151188  
 TITLE: Preparation of 3-phenyl-3,8-diazabicyclo[3.2.1]octanes and analogs as serotonin reuptake inhibitors  
 INVENTOR(S): Fliri, Anton Franz Josef; Gallaschun, Randall James  
 PATENT ASSIGNEE(S): Pfizer Products Inc., USA  
 SOURCE: Eur. Pat. Appl., 29 pp.  
 CODEN: EPXXDW  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1178047	A1	20020206	EP 2001-306313	20010723
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
BR 2001003180	A	20020326	BR 2001-3180	20010801
US 2002068748	A1	20020606	US 2001-920587	20010801
US 6531468	B2	20030311		
JP 2002088084	A2	20020327	JP 2001-235227	20010802
PRIORITY APPLN. INFO.: US 2000-222706P P 20000803				
OTHER SOURCE(S): CASREACT 136:151188; MARPAT 136:151188				




AB R3Zr1 [R1 = (un)substituted Ph; R3 = H, alkyl, (hetero)aryl, etc.; Z = e.g., 3,8-diazabicyclo[3.2.1]octane-8,3-diyl] were prepd. as serotonin reuptake inhibitors (no data). Thus, 1-benzyl-2,5-bis(chloromethyl)pyrrolidine (prepn. given) was cyclocondensed with 4-ClC6H4NH2 and the product hydrogenolized to give title compd. I.  
 IT 395059-29-7B, 8-Benzyl-3-(4-fluorophenyl)-3,8-diazabicyclo[3.2.1]octane 395059-41-3P, 8-Benzyl-3-(4-chlorophenyl)-3,8-diazabicyclo[3.2.1]octane 395059-55-9P, 3-Benzyl-8-(4-chlorophenyl)-3,8-diazabicyclo[3.2.1]octane  
 RL: PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)  
 (prepn. of 3-phenyl-3,8-diazabicyclo[3.2.1]octanes and analogs as serotonin reuptake inhibitors)  
 RN 395059-29-7 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 3-(4-fluorophenyl)-8-(phenylmethyl)- (9CI) (CA INDEX NAME)


L17 ANSWER 6 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 2001:629422 CAPLUS  
 DOCUMENT NUMBER: 136:200161  
 TITLE: 3,8-Diazabicyclo-[3.2.1]-octane derivatives as analogues of ambasilide, a Class III antiarrhythmic agent  
 AUTHOR(S): Villa, S.; Barlocco, D.; Cignarella, G.; Papp, G. J.; Balaci, B.; Takacs, J.; Varro, A.; Borosy, A.; Keszler, K.; Matyus, P.  
 CORPORATE SOURCE: Istituto di Chimica Farmaceutica, Università di Milano, Milan, 20131, Italy  
 SOURCE: European Journal of Medicinal Chemistry (2001), 36(6), 495-506  
 CODEN: EUMCA5; ISSN: 0223-5234  
 PUBLISHER: Editions Scientifiques et Médicales Elsevier  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB Ambasilide, a representative of Class III antiarrhythmics, was reported to prolong the cardiac action potential duration in the dog, with little or no effect on Ca and Na currents. A series of ambasilide analogs have been prepd. possessing the 3,8-diazabicyclo-[3.2.1]-octane moiety instead of the 3,7-diazabicyclo-[3.3.1]-nonane present in ambasilide. The compds. were tested by both in vitro extracellular electrophysiol. assays and by the conventional microelectrode technique. Most compds. tested lengthened the effective refractory period (ERP) with no change or only a slight increase on the impulse conduction time (ICT). Similarly some of the tested compds. lengthened the action potential duration (APD), a typical Class III feature, without exerting any significant effect on the maximal rate of depolarization, therefore apparently lacking Class I antiarrhythmic activity.


IT 401514-09-8P 401514-13-4P  
 RL: PAC (Pharmacological activity); PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent)  
 (prepn. and conformation energy anal. of antiarrhythmic diazabicyclooctanes as analogs of ambasilide)  
 RN 401514-09-8 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8-(4-aminobenzoyl)-3-(phenylmethyl)- (9CI) (CA INDEX NAME)




RN 401514-13-4 CAPLUS  
 CN Benzenamine, 4-[2-[3-(phenylmethyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]ethoxy]- (9CI) (CA INDEX NAME)




L17 ANSWER 5 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



RN 395059-41-3 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 3-(4-chlorophenyl)-8-(phenylmethyl)- (9CI) (CA INDEX NAME)



RN 395059-55-9 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8-(4-chlorophenyl)-3-(phenylmethyl)- (9CI) (CA INDEX NAME)




REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L17 ANSWER 6 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)

IT 401514-10-1P 401514-11-2P 401514-12-3P 401514-14-5P 401514-15-6P  
 RL: PAC (Pharmacological activity); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (prepn. and conformation energy anal. of antiarrhythmic diazabicyclooctanes as analogs of ambasilide)


RN 401514-10-1 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8-[4-[(methylsulfonyl)amino]benzoyl]-3-(phenylmethyl)- (9CI) (CA INDEX NAME)



RN 401514-11-2 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8-[4-[bis(methylsulfonyl)amino]benzoyl]-3-(phenylmethyl)- (9CI) (CA INDEX NAME)




RN 401514-12-3 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8-(4-nitrobenzoyl)-3-(phenylmethyl)- (9CI) (CA INDEX NAME)




RN 401514-14-5 CAPLUS  
 CN Acetamide, N-[4-[2-[3-(phenylmethyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]ethoxy]phenyl]- (9CI) (CA INDEX NAME)

09/972,177


L17 ANSWER 6 OF 39 CAPLUS COPYRIGHT 2003 ACS ON STN (Continued)



RN 401514-15-6 CAPLUS  
CN Methanesulfonamide, N-[4-[2-[3-(phenylmethyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]ethoxy]phenyl]- (9CI) (CA INDEX NAME)




IT 401514-18-9P  
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
(prepn. of antiarrhythmic diazabicyclooctanes via alkylation or amidation of N-protected diazabicyclooctane)  
RN 401514-18-9 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 3-(4-aminobenzoyl)-8-(phenylmethyl)- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 7 OF 39 CAPLUS COPYRIGHT 2003 ACS ON STN  
ACCESSION NUMBER: 2001:435076 CAPLUS  
DOCUMENT NUMBER: 135:46205  
TITLE: Preparation of neurotrophic bicyclic diamides with peptidylprolyl isomerase (PPIase or rotamase) inhibitory activity  
INVENTOR(S): Dubowchik, Gene Michael; Provencal, David Paul  
PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA  
SOURCE: PCT Int. Appl., 91 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001042245	A1	20010614	WO 2000-US32395	20001128
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
PRIORITY APPLN. INFO.: US 1999-169600P P 19991208				
OTHER SOURCE(S): MARPAT 135:46205				




AB The invention relates to the design, synthesis, and the peptidylprolyl isomerase (PPIase or rotamase) inhibitory activity of novel bicyclic diamide compds. that are neuroprotective and/or neurotrophic agents (i.e. compds. capable of stimulating growth or proliferation of nervous tissue), and that bind to immunophilins such as FKBP12 and inhibit their rotamase activity. This invention also relates to pharmaceutical compns. comprising these compds. The compds. are encompassed by structure I [X = O, F2; n = 1, 2; m = 0, 1, 2; p = 0, 1; D = alk(en)yl, cycloalk(en),

L17 ANSWER 7 OF 39 CAPLUS COPYRIGHT 2003 ACS ON STN (Continued)  
alk(en)yl, 2- or 3-indolyl, Ar, Ar-alk(en)yl; Ar = selected (un)substituted carbo- or heterocyclic arom. groups; Q, A = H, Ar, alk(en)yl, cycloalk(en)yl, their N/O/S-heteroat. analogs, etc.; and their pharmaceutically acceptable salts]. Over 40 examples were prepd. and tested. For instance, (1S,5R)-8-benzyl-3,8-diaza-3-(3-phenylpropyl)bicyclo[3.2.1]octan-2-one (prepn. given) underwent hydrogenolytic debenzoylation and amidation with 3,4,5-trimethoxyphenyl-2-oxoacetyl chloride to give title compd. II. In a fluorescence polarization assay of FKBP12 binding, II gave 34% inhibition at 1  $\mu$ M, and its 3-(3-pyridyloxy)propyl analog gave 98% inhibition.


IT 344462-40-4P 344462-41-5P 344462-42-6P  
344462-43-7P 344462-44-8P 344462-45-9P  
344462-47-1P 344462-48-2P 344462-49-3P  
344462-50-6P 344462-51-7P 344462-52-8P  
344462-53-9P 344462-55-1P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(stereoselective prepn. and biol. activity of bicyclic diamides as neuroprotective agents and peptidylprolyl isomerase (PPIase or rotamase) inhibitors)  
RN 344462-40-4 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 8-(phenylmethyl)-3-(3-phenylpropyl)-, (1S,5R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 344462-41-5 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 3-(4-phenylbutyl)-8-(phenylmethyl)-, (1S,5R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 344462-42-6 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 8-(phenylmethyl)-3-[3-(3,4,5-trimethoxyphenyl)propyl]-, (1S,5R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

L17 ANSWER 7 OF 39 CAPLUS COPYRIGHT 2003 ACS ON STN (Continued)




RN 344462-43-7 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 3-[2-(3,4-dimethoxyphenyl)ethyl]-8-(phenylmethyl)-, (1S,5R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 344462-44-8 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 3-[3-(3,4-dimethoxyphenyl)propyl]-8-(phenylmethyl)-, (1S,5R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 344462-45-9 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 3-[4-(3,4-dimethoxyphenyl)butyl]-8-(phenylmethyl)-, (1S,5R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

L17 ANSWER 7 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




RN 344462-47-1 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 8-(phenylmethyl)-3-[(3-phenyl-1-(2-phenylethyl)propyl)-, (1S,5R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 344462-48-2 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 3-[4-(3,4-dimethoxyphenyl)-1-(3-phenylpropyl)butyl]-8-(phenylmethyl)-, (1S,5R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 344462-49-3 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 8-(phenylmethyl)-3-[(2-(3-phenylpropyl)-5-(3,4,5-trimethoxyphenyl)pentyl)-, (1S,5R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

L17 ANSWER 7 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




RN 344462-50-6 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 8-(phenylmethyl)-3-[5-phenyl-2-(3-phenylpropyl)pentyl]-, (1S,5R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 344462-51-7 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 3-[5-(3,4-dimethoxyphenyl)-2-(3-(3,4-dimethoxyphenyl)propyl)pentyl]-8-(phenylmethyl)-, (1S,5R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 344462-52-8 CAPLUS

L17 ANSWER 7 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)


CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 3-[4-(3,4-dimethoxyphenyl)-2-[(3,4-dimethoxyphenyl)methyl]butyl]-8-(phenylmethyl)-, (1S,5R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 344462-53-9 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 3-[2-[2-(3,4-dimethoxyphenyl)ethyl]-4-phenylbutyl]-8-(phenylmethyl)-, (1S,5R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 344462-55-1 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 8-(phenylmethyl)-3-[(3-(3-pyridinyloxy)propyl)-, (1S,5R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

09/972,177


**L17 ANSWER 8 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN**  
 ACCESSION NUMBER: 2001:122158 CAPLUS  
 DOCUMENT NUMBER: 134:311179  
 TITLE: 3,8-Diazabicyclo[3.2.1]octan-2-one Peptide Mimetics: Synthesis of a Conformationally Restricted Inhibitor of Farnesyltransferase  
 AUTHOR(S): Dinsmore, Christopher J.; Bergman, Jeffrey M.; Bogusky, Michael J.; Culberson, J. Christopher; Hamilton, Kelly A.; Graham, Samuel L.  
 CORPORATE SOURCE: Departments of Medicinal Chemistry Molecular Systems and Cancer Research, Merck Research Laboratories, West Point, PA, 19486, USA  
 SOURCE: Organic Letters (2001), 3(6), 865-868  
 CODEN: ORLEF7; ISSN: 1523-7060  
 PUBLISHER: American Chemical Society  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 134:311179

**AB** A new synthesis of the 3,8-diazabicyclo[3.2.1]octan-2-one framework is described. Transannular enolate alkylation of piperazinone deriva. provides a flexible route to highly constrained bicyclic peptidomimetic synthons with substitution at the C.alpha. position. The chem. was used to produce a conformationally constrained farnesyltransferase inhibitor, which aided the elucidation of enzyme-bound conformation.

**IT** 335160-93-5P 335161-00-7P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (prepn. of a conformationally restricted farnesyltransferase inhibitor based on 3,8-diazabicyclo[3.2.1]octanone)



**RN** 335160-93-5 CAPLUS  
**CN** Benzonitrile, 4-[[5-[[[(1R,5S)-3-[(2,4-dimethoxyphenyl)methyl]-2-oxo-3,8-diazabicyclo[3.2.1]oct-8-yl)methyl]-1H-imidazol-1-yl)methyl]-2-fluoro-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



**RN** 335161-00-7 CAPLUS  
**CN** Benzonitrile, 4-[[5-[[[(1R,5S)-3-[(2-chloro-5-[(methylsulfonyl)oxy]phenyl)methyl]-2-oxo-3,8-diazabicyclo[3.2.1]oct-8-yl)methyl]-1H-imidazol-1-yl)methyl]-2-fluoro-(9CI) (CA INDEX NAME)

**L17 ANSWER 9 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN**  
 ACCESSION NUMBER: 1999:244249 CAPLUS  
 DOCUMENT NUMBER: 130:311770  
 TITLE: Benzocondensed derivatives as rigid analogs of the .mu.-opioid agonist 3(8)-cinnamyl-8(3)-propionyl-3,8-diazabicyclo[3.2.1]octanes: synthesis, modeling, and affinity  
 AUTHOR(S): Cignarella, G.; Barlocco, D.; Vianello, P.; Villa, S.; Pinna, G. A.; Fadda, P.; Fratta, W.; Toma, L.; Gessi, S.  
 CORPORATE SOURCE: Istituto di Chimica Farmaceutica e Tossicologica, Milan, 20131, Italy  
 SOURCE: Farmaco (1998), 53(10,11), 667-674  
 CODEN: FRMCE8; ISSN: 0014-827X  
 PUBLISHER: Elsevier Science S.A.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 GI




**AB** A new series of rigid analogs I and II (X = O, NH, S, CH=CH, CH2, (CH2)2, (CH2)3) of the previously reported analgesic 3-cinnamyl-8-propionyl-3,8-diazabicyclo[3.2.1]octane and its reverted isomer 3-propionyl-8-cinnamyl (III) were synthesized, in which the cinnamyl substituent is incorporated in benzocondensed bicyclic systems. Binding assays for the affinity towards .mu. receptors indicated that, while in the reverted series II the .beta.-naphthylmethyl and the benzocycloheptenylmethyl deriv. favorably compared with III, all compds. I displayed a .mu.-affinity lower than that of the parent. Modeling studies suggest that the flexibility of the cinnamyl side chain plays an important role for activity.

**IT** 172207-91-9P 223593-83-7P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (prepn. and .mu.-opioid agonist activity of cinnamylpropionyl-diazabicyclooctanes)


**RN** 172207-91-9 CAPLUS  
**CN** 3,8-Diazabicyclo[3.2.1]octane, 8-(1H-indol-2-ylcarbonyl)-3-(phenylmethyl)-(9CI) (CA INDEX NAME)

**L17 ANSWER 8 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN** (Continued)  
 Absolute stereochemistry.




**IT** 335160-83-3P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of a conformationally restricted farnesyltransferase inhibitor based on 3,8-diazabicyclo[3.2.1]octanone)  
**RN** 335160-83-3 CAPLUS  
**CN** 3,8-Diazabicyclo[3.2.1]octane-8-carboxylic acid, 3-[(2,4-dimethoxyphenyl)methyl]-2-oxo-1-(phenylmethyl)-, 1,1-dimethylethyl ester, (1S,5S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




**REFERENCE COUNT:** 39 THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

**L17 ANSWER 9 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN** (Continued)



**RN** 223593-83-7 CAPLUS  
**CN** 3,8-Diazabicyclo[3.2.1]octane, 8-(2-benzofuranylcarbonyl)-3-(phenylmethyl)-(9CI) (CA INDEX NAME)



**REFERENCE COUNT:** 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

09/972,177

L17 ANSWER 10 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN

ACCESSION NUMBER: 1998:88086 CAPLUS

DOCUMENT NUMBER: 128:97657

TITLE:

AUTHOR(S):

Mono- and Disubstituted-3,8-diazabicyclo[3.2.1]octane Derivatives as Analgesics Structurally Related to Epibatidine: Synthesis, Activity, and Modeling  
Barlocco, Daniela; Cignarella, Giorgio; Tondi, Donatella; Vianello, Paola; Villa, Stefania; Bartolini, Alessandro; Ghelardini, Carla; Galeotti, Nicoletta; Anderson, David J.; Kuntzweiler, Theresa A.; Colombo, Diego; Toma, Lucio  
Istituto di Chimica Farmaceutica e Tossico, Università Degli Studi di Milano, Milan, 20131, Italy  
Journal of Medicinal Chemistry (1998), 41(5), 674-681  
CODEN: JMCMAR; ISSN: 0022-2623

CORPORATE SOURCE:

SOURCE:

PUBLISHER:

DOCUMENT TYPE:

LANGUAGE:

AB A series of 3,8-diazabicyclo[3.2.1]octanes substituted either at the 3 position or at the 8 position by a chlorinated heteroaryl ring were synthesized, as potential analogs of the potent natural analgesic epibatidine. When tested in the hot plate assay, the majority of the compds. showed significant effects, the most interesting being the 3-(6-chloro-3-pyridazinyl)-3,8-diazabicyclo[3.2.1]octane (I). At a s.c. dose of 1 mg/kg, I induced a significant increase in the pain threshold, its action lasting for about 45 min. 1A also demonstrated good protection at a dose of 5 mg/kg in the mouse abdominal constriction test, while at 20 mg/kg it completely prevented the constrictions in the animals. Administration of naloxone (1 mg/kg i.p.) did not antagonize its antinociception while mecamylamine (2 mg/kg i.p.) did, thus suggesting the involvement of the nicotinic system in its action. Binding studies confirmed high affinity for the  $\alpha_4\beta_2$  nAChR subtype ( $K_i = 4.1 \pm 0.21$  nM). NACHR functional activity studies on three different cell lines showed that I was devoid of any activity at the neuromuscular junction. Finally, due to the analogy in their pharmacol. profile with that of epibatidine, compds. were compared from a structural and conformational point of view through theor. calcs. and high-field  $^1H$  NMR spectroscopy. Results indicate that all of them present one conformation similar to that of epibatidine.

IT 201162-43-8P


RI: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
(synthesis, activity, and modeling of mono- and disubstituted-3,8-diazabicyclo[3.2.1]octane derivs. as analgesics structurally related to epibatidine)

RN 201162-43-8 CAPLUS

CN 3,8-Diazabicyclo[3.2.1]octane, 8-(6-chloro-3-pyridazinyl)-3-(phenylmethyl)- (9CI) (CA INDEX NAME)

L17 ANSWER 10 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN

(Continued)



L17 ANSWER 11 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN

ACCESSION NUMBER: 1996:567069 CAPLUS

DOCUMENT NUMBER: 125:221856

TITLE:

INVENTOR(S):

Preparation of quinazoline derivatives as adrenergic  $\alpha_1$  receptor antagonists  
Andrews, Robert Carl; Brown, Peter Jonathan; Deaton, David Norman; Drewry, David Harold; Foley, Michael Andrew; Garrison, Deanna T.; Marron, Brian Edward; Smalley, Terrence L.; Berman, Judd M.; Noble, Stewart Alwyn

PATENT ASSIGNEE(S):

SOURCE:

Glaxo Inc, USA  
Brit. UK Pat. Appl., 190 pp.  
CODEN: BAXXDU

DOCUMENT TYPE:

LANGUAGE:

FAMILY ACC. NUM. COUNT: 1



PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
GB 2295387	A1	19960529	GB 1994-23635	19941123
PRIORITY APPLN. INFO.: GB 1994-23635 19941123				
OTHER SOURCE(S): MARPAT 125:221856				

GI

L17 ANSWER 11 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN

(Continued)



AB Title compds. [I: R = 2122 = R4; R1 = H, halo, alkyl, alkoxy, etc.; R4 = H, (di)alkyl amino, phenyl(omy), etc.; R5, R6 = H, OH, halo, alkyl, alkoxy; Z1 = NH, 2-(piperazine-1,4-diyl)ethylimino, iminopyridine-5,2-diylimino, etc.; Z2 = bond, (un)substituted alkylene] were prepd. as adrenergic  $\alpha_1$  receptor antagonists (no data). Thus, 4-chloro-2-phenylquinazoline was aminated by 4-amino-1-benzylpiperidine and the deprotected product N-alkylated by 5-(2-chloroethyl)-2-methoxybenzenesulfonamide (prepn. given) to give title compd. II.

IT 181115-69-5P

RI: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(prepn. of quinazoline derivs. as adrenergic  $\alpha_1$  receptor antagonists)


RN 181115-69-5 CAPLUS

CN 3,8-Diazabicyclo[3.2.1]octane, 3-(phenylmethyl)-8-(2-phenyl-4-quinazolinyl)- (9CI) (CA INDEX NAME)

09/972,177


L17 ANSWER 12 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1995:994354 CAPLUS  
 DOCUMENT NUMBER: 124:55984  
 TITLE: 3,8-Diazabicyclo[3.2.1]octane derivatives having analgesic activity  
 INVENTOR(S): Cignarella, Giorgio  
 PATENT ASSIGNEE(S): Riace Establishment, Liechtenstein  
 SOURCE: PCT Int. Appl., 21 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9523152	A1	19950831	WO 1995-EP476	19950210
W:	AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MX, NL, NO, NZ, PL, PT, RO, RU, SE, SI, SK, TJ, TT, UA, UG, US, UZ, VN			
RW:	KE, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
AU 9518085	A1	19950911	AU 1995-18085	19950210
EP 746560	A1	19961211	EP 1995-909700	19950210
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE			
US 5672601	A	19970930	US 1996-696948	19960822
PRIORITY APPLN. INFO.:			IT 1994-M1326	19940223
			WO 1995-EP476	19950210
OTHER SOURCE(S):			CASREACT 124:55984; MARPAT 124:55984	
GI				





AB Title compds. I and their pharmaceutically acceptable salts are claimed and/or prepd. [wherein R .noteq. R1; R, R1 = straight or branched C2-8 acyl, CH2AB; A = bond between 2 C atoms, CH2CH, or CH2CO; B = C6-10 aryl, (un)substituted with .gtoreq. 1 of CONHR, carboxy, cyano, NO2, or NHCOR; or (un)substituted arom. heterocyclic or alicyclic group with 5 or 6 members in the ring, optionally benzocondensed, contg. .gtoreq. 1 of N, O, or S; when R or R1 = propionyl, the other .noteq. cinnamyl or p-nitrocinnamyl; when R = propionyl, R1 .noteq. o- or m-nitrocinnamyl]. I have central analgesic activity comparable to morphine, and bind selectively to opioid .mu. receptors with similar affinity. However, I are substantially free of withdrawal phenomena, as detd. by the jumping test in mice, where activity was 3-20 times lower than morphine after 21 analgesically equipotent doses in 7 days (no addnl. data). For example,

L17 ANSWER 12 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
 N-alkylation of N8-acetyl-3,8-diazabicyclo[3.2.1]octane with cinnamyl chloride and K2CO3 in refluxing Me2CO gave I [R = Ac; R1 = CH2CH:CHPh].  
 IT 172207-91-99  
 RI: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (intermediate; prepn. of diazabicyclooctane derivs. as analgesics)  
 RN 172207-91-9 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8-(1H-indol-2-ylcarbonyl)-3-(phenylmethyl)- (9CI) (CA INDEX NAME)




L17 ANSWER 13 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1995:898142 CAPLUS  
 DOCUMENT NUMBER: 124:117785  
 TITLE: Concise synthesis of new homoaza sugars. Fully substituted, functionally diverse pyrrolidines  
 AUTHOR(S): Campanini, Laurence; Dureult, Annie; Depeyaz, Jean-Claude  
 CORPORATE SOURCE: Lab. Chim. Biochim. Pharmacologiques Toxicologiques, Univ. Rene Descartes, Paris, 75270, Fr.  
 SOURCE: Tetrahedron Letters (1995), 36(44), 8015-18  
 CODEN: TELEAY; ISSN: 0040-4039  
 PUBLISHER: Elsevier  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 GI




AB Five-membered deoxyaza sugars, e.g. I, of D-glucose configuration, bearing an aminomethyl, a bromomethyl or a thiomethyl group at the pseudo anomeric position, were prepd. by nucleophilic opening of C2 sym. bis-aziridines followed by chemoselective transformations of the nucleophile. The 1-bromo-2,5-imino-D-glucitol could be converted into attractive bicyclic compds., e.g. II.  
 IT 172795-11-89  
 RI: SPN (Synthetic preparation); PREP (Preparation)  
 (synthesis of pyrrolidine homoaza sugars via nucleophilic ring opening and intramol. cyclocondensation of bis-aziridines)  
 RN 172795-11-8 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 6,7-bis(phenylmethoxy)-3,8-bis(phenylmethyl)-, [1R-(6-endo,7-exo)]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L17 ANSWER 14 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1995:502804 CAPLUS  
 DOCUMENT NUMBER: 123:198662  
 TITLE: Synthesis of 5,7,8,9,10,11-hexahydro-7-oxo-8,11-iminoazepino[1,2-b]isoquinolines  
 AUTHOR(S): Peters, David A.; Yates, Nicholas D.; Scopes, David I. C.; Joule, John A.  
 CORPORATE SOURCE: Chem. Dep., Univ. Manchester, Manchester, M13 9PL, UK  
 SOURCE: Heterocycles (1995), 40(2), 983-91  
 CODEN: HETCYM; ISSN: 0385-5414  
 PUBLISHER: Japan Institute of Heterocyclic Chemistry  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 123:198662  
 AB Me 4-methylene-2-oxo-3,8-diazabicyclo[3.2.1]octane-6-exo-6-carboxylates have been converted into Me 5,7,8,9,10,11-hexahydro-7-oxo-8,11-iminoazepino[1,2-b]isoquinoline-10-carboxylates.  
 IT 167874-03-59  
 RI: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (synthesis of hexahydroiminoazepinoisoquinolines)  
 RN 167874-03-5 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-6-carboxylic acid, 3-[(2-bromophenyl)methyl]-4-methylene-2-oxo-8-(phenylmethyl)-, methyl ester, exo- (9CI) (CA INDEX NAME)

Relative stereochemistry.





L17 ANSWER 15 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1995:332276 CAPLUS  
 DOCUMENT NUMBER: 123:198733  
 TITLE: 1,3-Dipolar cycloadditions to oxidopyraziniums  
 AUTHOR(S): Yates, Nicholas D.; Peters, David A.; Allway, Philip A.; Beddoes, Roy L.; Scopes, David I. C.; Joule, John A.  
 CORPORATE SOURCE: Chem. Dep., Univ. Manchester, Manchester, M13 9PL, UK  
 SOURCE: Heterocycles (1995), 40(1), 331-47  
 CODEN: HTCYAM; ISSN: 0385-5414  
 PUBLISHER: Japan Institute of Heterocyclic Chemistry  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 GI




AB The cycloaddn. of dipolarophiles to oxidopyraziniums I (R1 = Me, R2 = Me, benzyl; R1 = 3-methoxybenzyl, R2 = Me) are described. Bicyclic products such as II are obtained.

IT 167418-00-0P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent) (prepn. of)

RN 167418-00-0 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 4-methylene-8-(phenylmethyl)-6-(phenylsulfonyl)-, exo- (9CI) (CA INDEX NAME)


Relative stereochemistry.




IT 167418-05-5P 167418-06-6P 167418-07-7P  
 167418-15-7P 167418-19-1P 167418-22-6P  
 RL: SPN (Synthetic preparation); PREP (Preparation)

RN 167418-05-5 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 4-methylene-8-(phenylmethyl)-6-(4-

L17 ANSWER 15 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




RN 167418-19-1 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 4-methoxy-4-methyl-8-(phenylmethyl)-6-(phenylsulfonyl)- (9CI) (CA INDEX NAME)




RN 167418-22-6 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-3-carboxylic acid, 4-methylene-2-oxo-8-(phenylmethyl)-6-(phenylsulfonyl)-, 1,1-dimethylethyl ester, exo- (9CI) (CA INDEX NAME)

Relative stereochemistry.




L17 ANSWER 15 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
 pyridinyl)-, endo- (9CI) (CA INDEX NAME)

Relative stereochemistry.




RN 167418-06-6 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 4-methylene-8-(phenylmethyl)-6-(2-pyridinyl)-, endo- (9CI) (CA INDEX NAME)

Relative stereochemistry.



RN 167418-07-7 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 4-methylene-6-phenyl-8-(phenylmethyl)-, endo- (9CI) (CA INDEX NAME)

Relative stereochemistry.



RN 167418-15-7 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 4-methyl-6-phenyl-8-(phenylmethyl)-, endo, endo- (9CI) (CA INDEX NAME)

Relative stereochemistry.

L17 ANSWER 16 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1994:77490 CAPLUS  
 DOCUMENT NUMBER: 120:77490  
 TITLE: The asymmetric synthesis of (-)-quinocarcin via a 1,3-dipolar cycloadditive strategy  
 AUTHOR(S): Garner, Philip; Ho, Wen Bin; Shin, Hunwoo  
 CORPORATE SOURCE: Dep. Chem., Case West. Reserve Univ., Cleveland, OH, 44106-7078, USA  
 SOURCE: Journal of the American Chemical Society (1993), 115(23), 10742-53  
 CODEN: JACSAT; ISSN: 0002-7863  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 120:77490  
 GI


\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB Details of the asym. synthesis and complete structure elucidation of (-)-quinocarcin (I), an antitumor antibiotic that inhibits DNA (and in some systems RNA) synthesis, are reported. Key steps in the synthesis include the use of an auxiliary-controlled 1,3-dipolar cycloaddn. reaction (II + III) (cf. IV) as well as an unprecedented intramol. imide olefination (V) (cf. VI) to assemble the 3,8-diazabicyclo[3.2.1]octane (CD ring) and isoquinoline (B ring) subunits of I in a stereo- and regiocontrolled manner. A comparison of the optical rotations of synthetic and natural quinocarcin confirms that the abs. configuration of this antibiotic is as depicted. Conclusive evidence for the (2aR) stereochem. in I is provided by a NOESY expt. on quinocarcin citrate.

IT 139527-59-6P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent) (prepn. and bromination of)


RN 139527-59-6 CAPLUS  
 CN 3H-3a,6-Methano-2,1-benzisothiazole, hexahydro-1-[[[3-[2-(methoxymethoxy)-1-(2-methoxy-6-methylphenyl)ethyl]-8-methyl-2,4-dioxo-3,8-diazabicyclo[3.2.1]oct-6-yl]carbonyl]-8,8-dimethyl-, 2,2-dioxide, {3aR-[1{1S\*,3(R\*),5R\*,6R\*},3a.alpha.,6.alpha.,7a.Beta.}]- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



IT 139527-61-0P


L17 ANSWER 16 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (prepn. and intramol. regioselective cyclization of)  
 RN 139527-61-0 CAPLUS  
 CN Phosphonium, [(3-methoxy-2-[2-(methoxymethoxy)-1-(8-methyl-2,4-dioxo-6-  
 (tetrahydro-8,8-dimethyl-2,2-dioxido-3H-3a,6-methano-2,1-benzisothiazol-  
 1(4H)-yl)carbonyl]-3,8-diazabicyclo[3.2.1]oct-3-  
 yl]ethyl)phenyl]methyl]triphenyl-, bromide, [3aR-  
 [1[1S\*,3(R\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX  
 NAME)



IT 139527-58-5P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (prepn. and methoxymethylation of)  
 RN 139527-58-5 CAPLUS  
 CN 3H-3a,6-Methano-2,1-benzisothiazole, hexahydro-1-[[3-(2-hydroxy-1-(2-  
 methoxy-6-methylphenyl)ethyl]-8-methyl-2,4-dioxo-3,8-  
 diazabicyclo[3.2.1]oct-6-yl]carbonyl]-8,8-dimethyl-, 2,2-dioxide,  
 [3aR-[1[1S\*,3(R\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX  
 NAME)

Absolute stereochemistry. Rotation (+).


AB ANSWER 17 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1993:551940 CAPLUS  
 DOCUMENT NUMBER: 119:151940  
 TITLE: Synthesis and opioid receptor affinity of bivalent  
 ligands derived from 3,8-diazabicyclo[3.2.1]octane  
 Barlocco, Daniela; Fadda, Paola; Fratta, Walter  
 CORPORATE SOURCE: Ist. Chim. Farm. Tozz., Univ. Milano, Milan, 20131,  
 Italy  
 SOURCE: Farmaco (1993), 48(3), 387-96  
 CODEN: FMCEB; ISSN: 0014-827X  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 GI



AB A new series of bivalent ligands [I, X = (CH2)2, (CH2)3, (CH2)4 or trans  
 CH2-CH=CH-CH2], derived from the previously reported analgesic  
 3-cinnamyl-8-propionyl-3,8-diazabicyclo[3.2.1]octane (II), has been  
 synthesized and tested in vitro for their affinity towards opioid  
 receptors and in vivo for their analgesic potency. None of the new  
 compds. showed either appreciable affinity for opioid receptors or  
 analgesic activity comparable to that of the model II.


IT 149771-39-1P 149771-40-4P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (prepn. and debenzilation of)  
 RN 149771-39-1 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8,8'-(1,4-dioxo-1,4-butanediyl)bis[3-  
 (phenylmethyl)- (9CI) (CA INDEX NAME)

L17 ANSWER 16 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)





IT 139527-60-9P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (prepn. and reaction of, with triphenylphosphine)  
 RN 139527-60-9 CAPLUS  
 CN 3H-3a,6-Methano-2,1-benzisothiazole, 1-[[3-[1-[2-(bromomethyl)-6-  
 methoxyphenyl]-2-(methoxymethoxy)ethyl]-8-methyl-2,4-dioxo-3,8-  
 diazabicyclo[3.2.1]oct-6-yl]carbonyl]hexahydro-8,8-dimethyl-, 2,2-dioxide,  
 [3aR-[1[1S\*,3(R\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX  
 NAME)


Absolute stereochemistry. Rotation (+).



L17 ANSWER 17 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)





RN 149771-40-4 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8,8'-(1,5-dioxo-1,5-pentanediy)bis[3-  
 (phenylmethyl)- (9CI) (CA INDEX NAME)



IT 149750-00-5P 150146-11-5P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and opioid receptor affinity of, analgesic activity in relation  
 to)  
 RN 149750-00-5 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8,8'-(1,4-dioxo-1,4-butanediyl)bis[3-  
 (phenylmethyl)-, dihydrochloride (9CI) (CA INDEX NAME)

09/972,177

L17 ANSWER 17 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)





● 2 HCl

RN 150146-11-5 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 8,8'-(1,5-dioxo-1,5-pentanediy)bis[3-(phenylmethyl)-, bis(4-methylbenzenesulfonate) (9CI) (CA INDEX NAME)




CM 1

CRN 149771-40-4  
CMF C31 H40 N4 O2



CM 2

~~L17 ANSWER 18 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN~~  
~~ACCESSION NUMBER: 1992:173839 CAPLUS~~  
~~DOCUMENT NUMBER: 116:173839~~  
~~TITLE: Asymmetric synthesis of (-)-quinocarcin~~  
~~AUTHOR(S): Garner, Philip Ho, Wen Bin; Shin, Hunwoo~~  
~~CORPORATE SOURCE: Dep. Chem., Case West. Reserve Univ., Cleveland, OH, 44106-7078, USA~~  
~~SOURCE: Journal of the American Chemical Society (1992), 114(7), 2767-8~~  
~~CODEN: JACSAT; ISSN: 0002-7863~~  
~~DOCUMENT TYPE: Journal~~  
~~LANGUAGE: English~~  
~~GI~~




AB The first asym. synthesis of (-)-quinocarcin (I) an antitumor antibiotic isolated from Streptomyces melanovineus that inhibits DNA (and in some systems RNA) synthesis, is reported. Key steps in the synthesis include an auxiliary-controlled 1,3-dipolar cycloaddn. reaction between imide II and acrylamide III and an unprecedented intramol. olefination of the imide IV to construct the 3,8-diazabicyclo[3.2.1]octane (CD ring) and isoquinoline (B-ring) subunits of I in a stereo- and regiocontrolled manner. A comparison of the optical rotations of synthetic and natural I confirms that the abs. configuration of this substance is as depicted.

IT 139527-59-6P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(prepn. and bromination of)  
RN 139527-59-6 CAPLUS  
CN 3H-3a,6-Methano-2,1-benzisothiazole, hexahydro-1-[[3-[2-(methoxymethoxy)-1-(2-methoxy-6-methylphenyl)ethyl]-8-methyl-2,4-dioxo-3,8-diazabicyclo[3.2.1]oct-6-yl]carbonyl]-8,8-dimethyl-, 2,2-dioxide, [3aR-[1[15\*,3(R\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX NAME)


Page 19

L17 ANSWER 17 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
CRN 104-15-4  
CMF C7 H8 O3 S




L17 ANSWER 18 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
NAME)


Absolute stereochemistry. Rotation (-).



IT 139527-61-0P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
(prepn. and cyclization of)  
RN 139527-61-0 CAPLUS  
CN Phosphonium, [[3-methoxy-2-[2-(methoxymethoxy)-1-[8-methyl-2,4-dioxo-6-[[tetrahydro-8,8-dimethyl-2,2-dioxido-3H-3a,6-methano-2,1-benzisothiazol-1(4H)-yl]carbonyl]-3,8-diazabicyclo[3.2.1]oct-3-yl]ethyl]phenyl]methyl]triphenyl-, bromide, [3aR-[1[15\*,3(R\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX NAME)




● 2 Br-



IT 139527-58-5P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)


L17 ANSWER 18 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
 (prepn. and etherification of)  
 RN 139527-58-5 CAPLUS  
 CN 3H-3a,6-Methano-2,1-benzisothiazole, hexahydro-1-[[3-[2-hydroxy-1-(2-methoxy-6-methylphenyl)ethyl]-8-methyl-2,4-dioxo-3,8-diazabicyclo[3.2.1]oct-6-yl]carbonyl]-8,8-dimethyl-, 2,2-dioxide, [3aR-[1[1S\*,3(R\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).




IT 139527-60-9P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (prepn. and reaction of, with triphenylphosphine)  
 RN 139527-60-9 CAPLUS  
 CN 3H-3a,6-Methano-2,1-benzisothiazole, 1-[[3-[1-[2-(bromomethyl)-6-methoxyphenyl]-2-(methoxymethoxy)ethyl]-8-methyl-2,4-dioxo-3,8-diazabicyclo[3.2.1]oct-6-yl]carbonyl]hexahydro-8,8-dimethyl-, 2,2-dioxide, [3aR-[1[1S\*,3(R\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (+).



L17 ANSWER 19 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
 (CA INDEX NAME)




IT 127381-65-1P 127470-56-8P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (prepn. and ethanolysis of)  
 RN 127381-65-1 CAPLUS  
 CN 3H-3a,6-Methano-2,1-benzisothiazole, 1-[[3-[2-[[[1,1-dimethylethyl]dimethylsilyl]oxy]-1-phenylethyl]-8-methyl-2,4-dioxo-3,8-diazabicyclo[3.2.1]oct-6-yl]carbonyl]hexahydro-8,8-dimethyl-, 2,2-dioxide, [3aS-[1[1S\*,3(R\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX NAME)



RN 127470-56-8 CAPLUS  
 CN 3H-3a,6-Methano-2,1-benzisothiazole, hexahydro-8,8-dimethyl-1-[[8-methyl-2,4-dioxo-3-(phenylmethyl)-3,8-diazabicyclo[3.2.1]oct-6-yl]carbonyl]-, 2,2-dioxide, [3aS-[1[1S\*,3(R\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX NAME)


L17 ANSWER 19 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1991:559499 CAPLUS  
 DOCUMENT NUMBER: 115:159499  
 TITLE: Development of an asymmetric approach to the 3,8-diazabicyclo[3.2.1]octane moiety of quinoxaline via intramolecular 1,3-dipolar cycloadditions of photochemically generated azomethine ylides  
 AUTHOR(S): Garner, Philip H.; Ho, Wen Bin; Grandhee, Sunitha K.; Youngs, Wiley J.; Kennedy, Vance O.  
 CORPORATE SOURCE: Dep. Chem., Case West. Reserve Univ., Cleveland, OH, 44106-7078, USA  
 SOURCE: Journal of Organic Chemistry (1991), 56(20), 5893-903  
 CODEN: JOCEAH; ISSN: 0022-3263  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 115:159499  
 GI




AB Exploratory work culminating in an enantioselective approach to the DNA-reactive alkaloid quinoxaline (I) is detailed. The key step involves auxiliary-controlled dipolar cycloaddn. between photochem. generated azomethine ylides such as II (R = H, CH<sub>2</sub>OSiMe<sub>2</sub>Me<sub>3</sub>) and Oppolzer's chiral acryloyl sultam (III) to assemble the 6-exo-substituted 3,8-diazabicyclo[3.2.1]octane core of I. The expected re-face selectivity of III was confirmed in one case by x-ray crystallog. anal. of endo-adduct. Removal (and recovery) of the chiral sultam auxiliary can be affected by titanium(IV)-mediated alcoholysis to give ester derivs. of the cycloadducts IV.

IT 127381-61-7P  
 RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and crystal structure of)  
 RN 127381-61-7 CAPLUS  
 CN 3H-3a,6-Methano-2,1-benzisothiazole, hexahydro-8,8-dimethyl-1-[[8-methyl-2,4-dioxo-3-(phenylmethyl)-3,8-diazabicyclo[3.2.1]oct-6-yl]carbonyl]-, 2,2-dioxide, [3aS-[1[1R\*,5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI)

L17 ANSWER 19 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




IT 127381-62-8P 127381-63-9P 127381-64-0P  
 127420-42-2P 127470-57-9P 135457-93-1P  
 135481-27-5P 135557-56-1P 135557-57-2P  
 135558-14-4P 135558-15-5P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of)  
 RN 127381-62-8 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-3-acetic acid, 8-methyl-2,4-dioxo-.alpha.-phenyl-6-[[[tetrahydro-8,8-dimethyl-2,2-dioxido-3H-3a,6-methano-2,1-benzisothiazol-1(4H)-yl]carbonyl]-, methyl ester, [3aS-[1[1S\*,3(S\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX NAME)




RN 127381-63-9 CAPLUS  
 CN 3H-3a,6-Methano-2,1-benzisothiazole, 1-[[3-[2-(acetyloxy)-1-phenylethyl]-8-methyl-2,4-dioxo-3,8-diazabicyclo[3.2.1]oct-6-yl]carbonyl]hexahydro-8,8-dimethyl-, 2,2-dioxide, [3aS-[1[1S\*,3(R\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX NAME)

L17 ANSWER 19 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



RN 127381-64-0 CAPLUS


CN 3H-3a,6-methano-2,1-benzisothiazole, 1-[[3-[2-(acetyloxy)-1-phenylethyl]-8-methyl-2,4-dioxo-3,8-diazabicyclo[3.2.1]oct-6-yl]carbonyl]hexahydro-8,8-dimethyl-2,2-dioxide, [3aR-[1[1S\*,3(S\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.eta.]]- (9CI) (CA INDEX NAME)



RN 127420-42-2 CAPLUS


CN 3,8-Diazabicyclo[3.2.1]octane-3-acetic acid, 8-methyl-2,4-dioxo-.alpha.-phenyl-6-[(tetrahydro-8,8-dimethyl-2,2-dioxido-3H-3a,6-methano-2,1-benzisothiazol-1(4H)-yl)carbonyl]-, methyl ester, [3aS-[1[1S\*,3(R\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX NAME)

L17 ANSWER 19 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



RN 127470-57-9 CAPLUS

CN 3H-3a,6-methano-2,1-benzisothiazole, 1-[[3-[2-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-1-phenylethyl]-8-methyl-2,4-dioxo-3,8-diazabicyclo[3.2.1]oct-6-yl]carbonyl]hexahydro-8,8-dimethyl-2,2-dioxide, [3aR-[1[1S\*,3(S\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX NAME)




RN 135457-93-1 CAPLUS

CN 3,8-Diazabicyclo[3.2.1]octane-6-carboxylic acid, 8-methyl-2,4-dioxo-3-(phenylmethyl)-, 5-methyl-2-(1-methylethyl)cyclohexyl ester, [1R-[1.alpha.,5.alpha.,6.alpha.(1R\*,2S\*,5R\*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


L17 ANSWER 19 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



RN 135481-27-5 CAPLUS

CN 3,8-Diazabicyclo[3.2.1]octane-6-carboxylic acid, 8-methyl-2,4-dioxo-3-(phenylmethyl)-, 1-[[[(dicyclohexylamino)sulfonyl]methyl]-7,7-dimethylbicyclo[2.2.1]hept-2-yl] ester, [1R-[1.alpha.,5.alpha.,6.alpha.(1S\*,2R\*,4R\*)]]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 135557-56-1 CAPLUS

CN 3,8-Diazabicyclo[3.2.1]octane-6-carboxylic acid, 8-methyl-2,4-dioxo-3-(phenylmethyl)-, 5-methyl-2-(1-methylethyl)cyclohexyl ester, [1S-[1.alpha.,5.alpha.,6.alpha.(1S\*,2R\*,5S\*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L17 ANSWER 19 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)

RN 135557-57-2 CAPLUS

CN 3,8-Diazabicyclo[3.2.1]octane-6-carboxylic acid, 8-methyl-2,4-dioxo-3-(phenylmethyl)-, 5-methyl-2-(1-methylethyl)cyclohexyl ester, [1R-[1.alpha.,5.alpha.,6.alpha.(1R\*,2S\*,5R\*)]]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 135558-14-4 CAPLUS

CN 3,8-Diazabicyclo[3.2.1]octane-6-carboxylic acid, 8-methyl-2,4-dioxo-3-(phenylmethyl)-, 5-methyl-2-(1-methylethyl)cyclohexyl ester, [1S-[1.alpha.,5.alpha.,6.alpha.(1S\*,2R\*,5S\*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 135558-15-5 CAPLUS


CN 3,8-Diazabicyclo[3.2.1]octane-6-carboxylic acid, 8-methyl-2,4-dioxo-3-(phenylmethyl)-, 1-[[[(dicyclohexylamino)sulfonyl]methyl]-7,7-dimethylbicyclo[2.2.1]hept-2-yl] ester, [1S-[1.alpha.,5.alpha.,6.alpha.(1R\*,2S\*,4S\*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L17 ANSWER 19 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




~~17~~ ANSWER 20 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 AB 1991:536722 CAPLUS  
 ACCESSION NUMBER: 115:136722  
 DOCUMENT NUMBER:  
 TITLE: Novel ring contractions via [2,3] Wittig type rearrangements: synthesis of 2-desoxy-2-methylenebicyclomycin  
 AUTHOR(S): Williams, Robert M.; Sabol, Mark R.; Kim, Hee Do; Kwast, Andrzej  
 CORPORATE SOURCE: Dep. Chem., Colorado State Univ., Fort Collins, CO, 80523, USA  
 SOURCE: Journal of the American Chemical Society (1991), 113(17), 6621-33  
 CODEN: JACSAT; ISSN: 0002-7863  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 OTHER SOURCE(S): CASREACT 115:136722  
 GI




AB Generation of bridgehead carbanions from bicyclo[5.2.2]- and bicyclo[7.2.2]-allyl ether-bridged piperazinediones results in novel ring contractions via unusual [2,3] Wittig and [3,3] Claisen rearrangements. The [2,3] Wittig rearrangement was applied to the oxadiazabicyclo[3.2.1]octane-2,4-dione I (R = 4-MeOC6H4CH2) in the construction of 2-desoxy-2-methylenebicyclomycin (II).


IT 135365-99-0P 135366-01-7P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of)  
 RN 135365-99-0 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-2,4-dione, 6-ethenyl-5-hydroxy-3,8-bis[(4-methoxyphenyl)methyl]-1-methyl- (9CI) (CA INDEX NAME)



L17 ANSWER 20 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
 RN 135366-01-7 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-2,4-dione, 1-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]-6-ethenyl-5-hydroxy-3,8-bis[(4-methoxyphenyl)methyl]- (9CI) (CA INDEX NAME)




~~17~~ ANSWER 21 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 AB 1990:424309 CAPLUS  
 ACCESSION NUMBER: 113:24309  
 DOCUMENT NUMBER:  
 TITLE: Stereoselective 1,3-dipolar cycloadditions of photochemically generated azomethine ylides to Oppolzer's chiral acryloyl sultam. An asymmetric approach to quinocarcin  
 AUTHOR(S): Garner, Philip; Ho, Wen Bin  
 CORPORATE SOURCE: Dep. Chem., Case West. Reserve Univ., Cleveland, OH, 44106-2699, USA  
 SOURCE: Journal of Organic Chemistry (1990), 55(13), 3973-5  
 CODEN: JOCEAH; ISSN: 0022-3263  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 GI




AB Photochem. generated azomethine ylides I [R = PhCH2, PhCH(CO2Me), PhCH(CH2OAc), PhCH(CH2OSiMe2OMe3)] underwent highly selective (ds >25:1) 1,3-dipolar cycloaddns. to the chiral acryloyl sultam (-)-II giving cycloadducts III corresponding to the substituted 3,8-diazabicyclo[3.2.1]octane moiety of quinocarcin with complete stereocontrol. The analogous reaction of with (+)-II provided the diastereomeric adducts, confirming that the stereochem. outcome is under control of the chiral auxiliary. The sultam auxiliary was readily removed (in recoverable form) by Ti(IV) -promoted alcoholysis of the cycloadducts.

IT 127381-62-8P 127381-65-1P 127420-42-2P  
 127470-56-8P 127470-57-9P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (prepn. and alcoholysis of)  
 RN 127381-62-8 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-3-acetic acid, 8-methyl-2,4-dioxo-.alpha.-phenyl-6-[(tetrahydro-8,8-dimethyl-2,2-dioxido-3H-3a,6-methano-2,1-benzisothiazol-1(4H)-yl)carbonyl]-, methyl ester, [3aS-[1[1S',3(S'),5R',6R'],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX NAME)

L17 ANSWER 21 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




RN 127381-65-1 CAPLUS  
 CN 3H-3a,6-Methano-2,1-benzisothiazole, 1-[[3-[2-[[[1,1-dimethylethyl]dimethylsilyl]oxy]-1-phenylethyl]-8-methyl-2,4-dioxo-3,8-diazabicyclo[3.2.1]oct-6-yl]carbonyl]hexahydro-8,8-dimethyl-, 2,2-dioxide, [3aS-[1[1S\*,3(R\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX NAME)




RN 127420-42-2 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-3-acetic acid, 8-methyl-2,4-dioxo-.alpha.-phenyl-6-[(tetrahydro-8,8-dimethyl-2,2-dioxido-3H-3a,6-methano-2,1-benzisothiazol-1(4H)-yl)carbonyl]-, methyl ester, [3aS-[1[1S\*,3(R\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX NAME)

L17 ANSWER 21 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




RN 127470-56-8 CAPLUS  
 CN 3H-3a,6-Methano-2,1-benzisothiazole, hexahydro-8,8-dimethyl-1-[[8-methyl-2,4-dioxo-3-(phenylmethyl)-3,8-diazabicyclo[3.2.1]oct-6-yl]carbonyl]-, 2,2-dioxide, [3aS-[1[1S\*,3(R\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX NAME)




RN 127470-57-9 CAPLUS  
 CN 3H-3a,6-Methano-2,1-benzisothiazole, 1-[[3-[2-[[[1,1-dimethylethyl]dimethylsilyl]oxy]-1-phenylethyl]-8-methyl-2,4-dioxo-3,8-diazabicyclo[3.2.1]oct-6-yl]carbonyl]hexahydro-8,8-dimethyl-, 2,2-dioxide, [3aR-[1[1S\*,3(S\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX NAME)

L17 ANSWER 21 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




IT 127381-61-7P 127381-63-9P 127381-64-0P  
 RL: SPN (Synthetic preparation); PREP (Preparation) (prepn. of)  
 RN 127381-61-7 CAPLUS  
 CN 3H-3a,6-Methano-2,1-benzisothiazole, hexahydro-8,8-dimethyl-1-[[8-methyl-2,4-dioxo-3-(phenylmethyl)-3,8-diazabicyclo[3.2.1]oct-6-yl]carbonyl]-, 2,2-dioxide, [3aS-[1[1R\*,3(R\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX NAME)




RN 127381-63-9 CAPLUS  
 CN 3H-3a,6-Methano-2,1-benzisothiazole, 1-[[3-[2-(acetyloxy)-1-phenylethyl]-8-methyl-2,4-dioxo-3,8-diazabicyclo[3.2.1]oct-6-yl]carbonyl]hexahydro-8,8-dimethyl-, 2,2-dioxide, [3aS-[1[1S\*,3(R\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX NAME)

L17 ANSWER 21 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



RN 127381-64-0 CAPLUS  
 CN 3H-3a,6-Methano-2,1-benzisothiazole, 1-[[3-[2-(acetyloxy)-1-phenylethyl]-8-methyl-2,4-dioxo-3,8-diazabicyclo[3.2.1]oct-6-yl]carbonyl]hexahydro-8,8-dimethyl-, 2,2-dioxide, [3aR-[1[1S\*,3(S\*),5R\*,6R\*],3a.alpha.,6.alpha.,7a.beta.]]- (9CI) (CA INDEX NAME)




09/972,177


D17 ANSWER 22 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1989:57646 CAPLUS  
 DOCUMENT NUMBER: 110:57646  
 TITLE: Antibacterial naphthyridine- and quinolonecarboxylic acid derivatives  
 INVENTOR(S): Weber, Abraham; Bouzard, Daniel; Essiz, Munir; Di Cesare, Pierre; Jacquet, Jean Pierre; Remuzon, Phillippe  
 PATENT ASSIGNEE(S): Bristol-Myers Co., USA  
 SOURCE: PCT Int. Appl., 100 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 8802627	A1	19880421	WO 1987-US2556	19871008
W: AU, DK, FI, HU, JP, KR, NO, RO, US				
RW: AT, BE, CH, DE, FR, GB, IT, LU, NL, SE				
ZA 8707471	A	19880525	ZA 1987-7471	19871005
DD 266354	A5	19890329	DD 1987-307706	19871006
DD 280530	A5	19900711	DD 1987-327989	19871006
AU 8781581	A1	19880506	AU 1987-81581	19871008
AU 611400	B2	19910613		
EP 288519	A1	19881102	EP 1987-907178	19871008
R: AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE				
HU 52500	A2	19900728	HU 1986-56	19871008
HU 203753	B	19910930		
DK 8803555	A	19880823	DK 1988-3555	19880628
NO 8803077	A	19880822	NO 1988-3077	19880708
FI 8803894	A	19880823	FI 1988-3894	19880823
CS 270598	B2	19900712	CS 1988-7400	19881110
AU 9176326	A1	19910808	AU 1991-76326	19910501
PRIORITY APPLN. INFO.:			US 1986-916752	19861008
			CS 1987-7295	19871008
			WO 1987-US2556	19871008
OTHER SOURCE(S):		MARPAT 110:57646		
GI				

L17 ANSWER 22 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




AB The title compds. I [X = F, Cl, Br, CF<sub>3</sub>, CCl<sub>3</sub>; Z = Q1, Q2, etc.; A, B, C, D, = H, (substituted) lower alkyl, NH<sub>2</sub>, OH, F, Cl, etc.; n = 0-3; R1 = CH<sub>3</sub>, CH<sub>2</sub>CH<sub>2</sub>Me, CPhMe<sub>2</sub>, etc.; R2 = H, C1-4 alkyl, alkali and alk. earth metal ions; R3 = H, (substituted) C1-6 alkyl, C3-6 cycloalkyl, etc.; Y = CH, CF, CCl, CBr, N], useful as antibacterials, were prepd., e.g., using amines II, III, IV, etc. Reaction of Et 1-(1,1-dimethylethyl)-1,4-dihydro-6,7,8-trifluoro-4-oxo-3-quinolinecarboxylate with piperazine in MeCN, followed by sapon. and workup, gave 7-piperazinyl-1-(1,1-dimethylethyl)-1,4-dihydro-6,8-difluoro-4-oxo-3-quinolinecarboxylic acid (V). V in vitro exhibited a MIC of 4 .mu.g/mL against Pseudomonas aeruginosa. The corresponding MIC of norfloxacin was 0.5 .mu.g/mL.  
 IT 118329-60-5  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (reaction of, in prepn. of naphthyridine and quinolone antibacterials)  
 RN 118329-60-5 CAPLUS  
 CN 1,8-Naphthyridine-3-carboxylic acid, 1-(1,1-dimethylethyl)-6-fluoro-1,4-dihydro-4-oxo-7-[3-(phenylmethyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]-, ethyl ester (9CI) (CA INDEX NAME)




L17 ANSWER 22 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)

D17 ANSWER 23 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1985:488088 CAPLUS  
 DOCUMENT NUMBER: 103:88088  
 TITLE: Synthesis and pharmacological activity of 3-aminopropiophenones and 3-(aminomethyl)camphors  
 AUTHOR(S): Occelli, E.; Fontanella, L.; Diena, A.; Schiatti, P.  
 CORPORATE SOURCE: Lab. Ric., Gruppo Lepetit S.p.A., Milan, Italy  
 SOURCE: Farmaco, Edizione Scientifica (1985), 40(2), 86-101  
 CODEN: FRPSAX; ISSN: 0430-0920  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Italian  
 GI



AB Z:CPhCH<sub>2</sub>CH<sub>2</sub>NR<sub>2</sub>.HCl (I, R = alkyl, (substituted) N-contg. heterocyclyl; Z = O, (acyl) hydroxyimino] and the camphor deriva. II (R same as above) were prepd. and their CNS, analgesic, and antiinflammatory activities evaluated.  
 IT 97669-75-5P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and CNS activity of)  
 RN 97669-75-5 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-3-propanimine, N-[[[4-(methoxyphenyl)amino]carbonyloxy]-8-(2-nitrobenzoyl)-.alpha.-phenyl]- (9CI) (CA INDEX NAME)




IT 97670-11-6P  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (prepn. and pharmacol. activities of)  
 RN 97670-11-6 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-3-propanimine, N-[[[4-(methoxyphenyl)amino]carbonyloxy]-8-(2-nitrobenzoyl)-.alpha.-phenyl]-, monohydrochloride (9CI) (CA INDEX NAME)




09/972,177

L17 ANSWER 23 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



● HCl

IT 97669-87-9P 97669-99-3P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of)  
 RN 97669-87-9 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8-(2-nitrobenzoyl)-3-(3-oxo-3-phenylpropyl)-, monohydrochloride (9CI) (CA INDEX NAME)




● HCl

RN 97669-99-3 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-3-propanimine, N-hydroxy-8-(2-nitrobenzoyl)-.alpha.-phenyl-, ethanedioate (1:1) (salt) (9CI) (CA INDEX NAME)

CM 1

CRN 97669-98-2  
 CMF C22 H24 N4 O4




CM 2

CRN 144-62-7  
 CMF C2 H2 O4

L17 ANSWER 24 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1984:423500 CAPLUS  
 DOCUMENT NUMBER: 101:23500  
 TITLE: Diazabicyclooctanes with anxiolytic and sedative activity  
 INVENTOR(S): Pedrazzoli, Andrea; Crisafulli, Emilio  
 PATENT ASSIGNEE(S): Sanofi, Fr.  
 SOURCE: Fr. Demande, 14 pp.  
 CODEN: FRXXBL  
 DOCUMENT TYPE: Patent  
 LANGUAGE: French  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 2531709	A1	19840217	FR 1982-14127	19820813
FR 2531709	B1	19850111		

PRIORITY APPL. INFO.: FR 1982-14127 19820813  
 OTHER SOURCE(S): CASREACT 101:23500  
 GI



AB 3,8-Diazabicyclo[3.2.1]octanes I [one of R and R1 is 2-pyrimidinyl and the other is H, alkyl, Ph, tolyl, PhCH2, 3,4-(CH2O2)C6H3CH2, PhCH:CHCH2, alkanoyl, PhCO, 3,4-(CH2O2)C6H2CO, PhCH:CHCO], which were prepd., are useful as anxiolytics and sedatives (no data). I [R = 3,4-(CH2O2)C6H3CO, R1 = H] was treated with 2-chloropyrimidine and K2CO3 in DMF to give I [R = 3,4-(CH2O2)C6H3CO, R1 = 2-pyrimidinyl].

IT 90478-34-5P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (prepn. and hydride redn. of)  
 RN 90478-34-5 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-2,4-dione, 3-(2-methylphenyl)-8-(phenylmethyl)-, monohydrochloride (9CI) (CA INDEX NAME)




● HCl

IT 90478-31-2P 90478-35-6P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)


L17 ANSWER 23 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




L17 ANSWER 24 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
 (prepn. and hydrogenolysis of)  
 RN 90478-31-2 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8-(1,3-benzodioxol-5-ylcarbonyl)-3-(phenylmethyl)- (9CI) (CA INDEX NAME)



RN 90478-35-6 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 3-(2-methylphenyl)-8-(phenylmethyl)- (9CI) (CA INDEX NAME)




IT 90478-39-0 90478-40-3 90478-41-4  
 90478-45-8 90478-51-6 90478-52-7  
 90478-53-8  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (N-alkylation by, of diazabicyclooctane deriv.)  
 RN 90478-39-0 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8-(1,3-benzodioxol-5-ylcarbonyl)-3-(2-pyrimidinyl)- (9CI) (CA INDEX NAME)




RN 90478-40-3 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8-(1,3-benzodioxol-5-ylmethyl)-3-(2-pyrimidinyl)- (9CI) (CA INDEX NAME)

L17 ANSWER 24 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



RN 90478-41-8 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 8-(1,3-benzodioxol-5-ylmethyl)-3-(2-pyrimidinyl)-, dihydrochloride (9CI) (CA INDEX NAME)



● 2 HCl

RN 90478-45-8 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 3-(phenylmethyl)-8-(2-pyrimidinyl)- (9CI) (CA INDEX NAME)

L17 ANSWER 25 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN

ACCESSION NUMBER: 1979:72149 CAPLUS

DOCUMENT NUMBER: 90:72149

TITLE: Tricyclic homologs of piperazine. III. Synthesis of 4-substituted hexahydro-1H-2,6-methanopyrrolo[1,2-a]pyrazines

AUTHOR(S): Occelli, E.; Fontanella, L.; Testa, E.

CORPORATE SOURCE: Lab. Ric., Lepetit S.p.A., Milan, Italy


SOURCE: Farmaco, Edizione Scientifica (1978), 33(11), 875-84

CODEN: FRPSAX; ISSN: 0430-0920

DOCUMENT TYPE: Journal

LANGUAGE: Italian

GI




AB Methanopyrrolopyrazine I (R = H, R1 = R2 = Me, X = Cl) was prepd. by treating II (R3 = H, R4 = CH2Ph) with BrCHMeCO2Et, debenzylating II (R3 = CHMeCO2Et, R4 = CH2Ph), methylating II (R3 = CHMeCO2Et, R4 = H), reducing II (R3 = CHMeCO2Et, R4 = Me), chlorinating the resulting alc., and cyclizing II (R3 = CHMeCH2Cl, R4 = Me) with base. I (R = R1 = Me, R2 = CH2Ph, X = MeSO3; R = H, R1 = Ph, R2 = CH2Ph, X = Cl) were similarly prepd. The latter 2 compds. gave 23 and 64 decrease resp. in gastrocnemius muscle contraction in rabbits at 2 mg/kg i.v.

IT 69099-93-0P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent) (prepn. and chlorination of)

RN 69099-93-0 CAPLUS

CN 3,8-Diazabicyclo[3.2.1]octane-8-ethanol, .beta.-phenyl-3-(phenylmethyl)-, dihydrochloride (9CI) (CA INDEX NAME)



● 2 HCl


IT 69099-94-1P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent) (prepn. and cyclization of)


RN 69099-94-1 CAPLUS

CN 3,8-Diazabicyclo[3.2.1]octane, 8-(2-chloro-1-phenylethyl)-3-(phenylmethyl)-, dihydrochloride (9CI) (CA INDEX NAME)


L17 ANSWER 24 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




RN 90478-51-6 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 3-(1,3-benzodioxol-5-ylcarbonyl)-8-(2-pyrimidinyl)- (9CI) (CA INDEX NAME)



RN 90478-52-7 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 3-(1,3-benzodioxol-5-ylmethyl)-8-(2-pyrimidinyl)- (9CI) (CA INDEX NAME)




RN 90478-53-8 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 3-(1,3-benzodioxol-5-ylmethyl)-8-(2-pyrimidinyl)-, dihydrochloride (9CI) (CA INDEX NAME)



● 2 HCl

L17 ANSWER 25 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




● 2 HCl

IT 69099-91-8P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent) (prepn. and redn. of)

RN 69099-91-8 CAPLUS

CN 3,8-Diazabicyclo[3.2.1]octane-8-acetic acid, .alpha.-phenyl-3-(phenylmethyl)-, methyl ester (9CI) (CA INDEX NAME)




IT 69099-92-9P

RL: SPN (Synthetic preparation); PREP (Preparation) (prepn. of)

RN 69099-92-9 CAPLUS

CN 3,8-Diazabicyclo[3.2.1]octane-8-ethanol, .beta.-phenyl-3-(phenylmethyl)-, dihydrochloride (9CI) (CA INDEX NAME)




09/972,177

L<sup>1</sup> ANSWER 26 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1978:509371 CAPLUS  
 DOCUMENT NUMBER: 89:109371  
 TITLE: Bicyclic piperazine homologs. XIV. Synthesis and analgesic activity of 3,8-diazabicyclo[3.2.1]octane derivatives  
 AUTHOR(S): Occeili, E.; Fontanella, L.; Diana, A.  
 CORPORATE SOURCE: Lab. Ric., Gruppo Lepetit S.p.A., Milan, Italy  
 SOURCE: Farmaco, Edizione Scientifica (1978), 33(6), 401-20  
 CODEN: FRPSAX; ISSN: 0430-0920  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Italian  
 GI



AB 2,6-Ethanopiperazines I [R = CH<sub>2</sub>CH:Me<sub>2</sub>, PrCO, EtCO; R<sub>1</sub> = CH<sub>2</sub>CH:Me<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>C(OR<sub>2</sub>)Me<sub>2</sub> (R<sub>2</sub> = H, acyl)] were prepd. by known methods and exhibited analgesic activity.  
 IT 67572-27-4P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (prepn. and hydrogenolysis of)  
 RN 67572-27-4 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8-benzoyl-3-(phenylmethyl)-, monohydrochloride (9CI) (CA INDEX NAME)




● HCl

L<sup>1</sup> ANSWER 28 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1972:539972 CAPLUS  
 DOCUMENT NUMBER: 77:139972  
 TITLE: Bicyclic analogs of piperazine. X. N-amino derivatives of 3,8-diazabicyclo[3.2.1]octane, 3,8-diazabicyclo[3.2.1]octane-2,4-diones, and 2,6-dimethylpiperazine with potential pharmacological activity  
 AUTHOR(S): Fontanella, L.; Occeili, E.; Testa, E.; Cignarella, G.  
 CORPORATE SOURCE: Lab. Ric., Gruppo Lepetit S.p.A., Milan, Italy  
 SOURCE: Farmaco, Edizione Scientifica (1972), 27(9), 755-72  
 CODEN: FRPSAX; ISSN: 0430-0920  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Italian  
 GI For diagram(s), see printed CA Issue.


AB The diazabicyclooctanes I [R = NO, H, Me, CH<sub>2</sub>, Ph, CH<sub>2</sub>CH:CHPh, COEt, NH<sub>2</sub>; R<sub>1</sub> = Me, COEt, CO<sub>2</sub>Et, NO, NH<sub>2</sub>, NHCOC<sub>6</sub>H<sub>3</sub>(SO<sub>2</sub>NH<sub>2</sub>)Cl-3,4, the diazabicyclooctane-diones II (R = NO, NH<sub>2</sub>, substituted amino) and some related piperazine derivs. were prepd. for testing for pharmacol. activity. I (R = NHCOC<sub>2</sub>Et, R<sub>1</sub> = CH<sub>2</sub>CH:CHPh) had anticonvulsant, analgesic, and local anesthetic activity and I (R = COEt, R<sub>2</sub> = Me, CH<sub>2</sub>CH:CHPh) also showed some activity. II (R = 3,4-(MeO)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>CH<sub>2</sub>NH) had slight analgesic activity. The piperazines had considerably lower diuretic activity than Clonamide.

IT 38074-18-9P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of)  
 RN 38074-18-9 CAPLUS  
 CN Benzamide, 3-(aminosulfonyl)-4-chloro-N-[3-(phenylmethyl)-3,8-diazabicyclo[3.2.1]oct-8-yl]- (9CI) (CA INDEX NAME)




L<sup>1</sup> ANSWER 27 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1978:593231 CAPLUS  
 DOCUMENT NUMBER: 83:193231  
 TITLE: Bicyclic analogs of piperazine. XIII. Derivatives of 3,8-diazabicyclo[3.2.1]octanes with potential antiinflammatory activity  
 AUTHOR(S): Fontanella, L.; Occeili, E.; Testa, E.  
 CORPORATE SOURCE: Lab. Ric., Gruppo Lepetit S.p.A., Milan, Italy  
 SOURCE: Farmaco, Edizione Scientifica (1975), 30(9), 742-53  
 CODEN: FRPSAX; ISSN: 0430-0920  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Italian  
 GI For diagram(s), see printed CA Issue.


AB Diazabicyclooctanes I (R = Me, R<sub>2</sub> = acyl; R = carboxyalkyl, acyl, R<sub>1</sub> = Me; R = CO<sub>2</sub>Et, R<sub>1</sub> = substituted alkyl, isonicotinoyl; R = H, R<sub>1</sub> = cyclopropylmethyl; R = CH<sub>2</sub>CO<sub>2</sub>H, R<sub>1</sub> = COC<sub>6</sub>H<sub>4</sub>Cl-4) (40 compds.) were prepd. by acylation of I (R or R<sub>1</sub> = Me) reaction with alkyl chloride, or acylation and redn.  
 IT 57269-57-5P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (prepn. and debenzoylation of)  
 RN 57269-57-5 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 3-(cyclopropylmethyl)-8-(phenylmethyl)- (9CI) (CA INDEX NAME)



IT 57269-56-4P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (prepn. and redn. of)  
 RN 57269-56-4 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 8-benzoyl-3-(cyclopropylcarbonyl)- (9CI) (CA INDEX NAME)




L<sup>1</sup> ANSWER 29 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1972:514359 CAPLUS  
 DOCUMENT NUMBER: 77:114359  
 TITLE: Synthesis of N,N'-dibenzylpyrrolidine-2,5-dicarboximide (3,8-diazabicyclo[3.2.1]octane-2,4-dione)  
 AUTHOR(S): Della, E. W.; Kendall, M.  
 CORPORATE SOURCE: Sch. Phys. Sci., Flinders Univ. South Australia, Bedford Park, Australia  
 SOURCE: Australian Journal of Chemistry (1972), 25(8), 1827-8  
 CODEN: AJCHAS; ISSN: 0004-9425  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 GI For diagram(s), see printed CA Issue.  
 AB The title compd. (I) was prepd. by cyclizing the pyrrolidine (II, R = H, R<sub>1</sub> = PhCH<sub>2</sub>NH) (III), which was obtained by hydrolysis of the ester prepd. from II (R = Et, R<sub>1</sub> = OEt) by the method of S. W. Blackman and R. J. Baltzly (1961). Thus, III was treated with SO<sub>2</sub>Cl<sub>2</sub> to give 75% I.HCl.  
 IT 17740-41-9P 37061-44-2P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of)  
 RN 17740-41-9 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-2,4-dione, 3,8-bis(phenylmethyl)-, monohydrochloride (9CI) (CA INDEX NAME)




● HCl

RN 37061-44-2 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-2,4-dione, 3,8-bis(phenylmethyl)- (9CI) (CA INDEX NAME)




09/972,177

~~L17~~ ANSWER 30 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1972:85788 CAPLUS  
 DOCUMENT NUMBER: 76:85788  
 TITLE: Bicyclic homologs of piperazine. XI.  
 3,8-Diazabicyclo[3.2.1]octane-2,4-diones with  
 potential pharmacological activity  
 Fontanella, L.; Occelli, E.  
 AUTHOR(S): Lab. Ric., Gruppo Lepetit S.p.A., Milan, Italy  
 CORPORATE SOURCE: Farmaco, Edizione Scientifica (1972), 27(1), 68-78  
 SOURCE: CODEN: FRPSAX; ISSN: 0430-0920  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Italian  
 GI For diagram(s), see printed CA Issue.  
 AB The 3-substituted 3,8-diazabicyclo[3.2.1]octane-2,4-diones, I (R = H), are  
 alkylated and acylated and treated with isocyanates to give  
 3,8-disubstituted compds. I (R = H, R1 = Me) is treated with BuI to give  
 1 (R = Bu, R1 = Me). Similarly prepd. are .apprx.30 addnl. I (R = alkyl,  
 acyl, CONH2, CONHPh; R1 = H, Me, PhCH2, aryl). II is treated with NH3 to  
 give 1 (R = Me, R1 = H); and 1 (R = H, R1 = p-tolyl) is prepd. by the  
 distn. of III.  
 IT 35101-50-9 35101-51-0 35101-52-1  
 RL: PROC (Process)  
 (prepn. of)  
 RN 35101-50-9 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-2,4-dione, 8-benzoyl-3-(phenylmethyl)- (9CI) (CA INDEX NAME)




RN 35101-51-0 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-2,4-dione, 8-(phenylacetyl)-3-(phenylmethyl)- (9CI) (CA INDEX NAME)




RN 35101-52-1 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-2,4-dione, 8-(1-oxo-2-phenyl-2-propenyl)-3-(phenylmethyl)- (9CI) (CA INDEX NAME)


~~L17~~ ANSWER 31 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1972:3795 CAPLUS  
 DOCUMENT NUMBER: 76:3795  
 TITLE: Synthesis of 2,5- and 2,6-bis(bromomethyl)-1,4-diphenylpiperazines and their conversion into 2,5-diphenyl-2,5-diazabicyclo[2.2.2]octane  
 Nelson, David A.; Worman, James J.; Keen, Brian  
 AUTHOR(S): Dep. Chem., Univ. Wyoming, Laramie, WY, USA  
 CORPORATE SOURCE: Journal of Organic Chemistry (1971), 36(22), 3361-5  
 SOURCE: CODEN: JOCEAH; ISSN: 0022-3263  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 GI For diagram(s), see printed CA Issue.  
 AB Treatment of cis-1,5-diphenyl-3,7-dihydroxyoctahydro-1,5-diazocine with PBr3 yielded a mixt. of cis-2,6-bis(bromomethyl)-1,4-diphenylpiperazine (I) and cis-2,5-bis(bromomethyl)-1,4-diphenylpiperazine (II). The structures of I and II were confirmed by conversion to the corresponding dimethyl-1,4-diphenylpiperazines (III and IV) by LiAlH4. III and IV were synthesized from cis-2,5- and cis-2,6-dimethylpiperazines. Both I and II on treatment with Mg in THF were converted to 2,5-diphenyl-2,5-diazabicyclo[2.2.2]octane (V). The interconversion of I and II is discussed.  
 IT 17740-42-0  
 RL: PRP (Properties)  
 (nuclear magnetic resonance spectrum of)  
 RN 17740-42-0 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 3,8-bis(phenylmethyl)- (9CI) (CA INDEX NAME)




L17 ANSWER 30 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



IT 35142-72-4P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of)  
 RN 35142-72-4 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-2,4-dione, 8-benzoyl-3-phenyl- (9CI) (CA INDEX NAME)




~~L17~~ ANSWER 32 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1970:78780 CAPLUS  
 DOCUMENT NUMBER: 72:78780  
 TITLE: Photoinduced 1,3-dipolar cycloaddition reaction of aziridinedicarboximide  
 Oida, Sadao; Ohki, Eiji  
 AUTHOR(S): Cent. Res. Lab., Sankyo Co., Ltd., Tokyo, Japan  
 CORPORATE SOURCE: Chemical & Pharmaceutical Bulletin (1969), 17(12), 2461-74  
 SOURCE: CODEN: CPBTAL; ISSN: 0009-2363  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 GI For diagram(s), see printed CA Issue.  
 AB Cycloaddn. of MeO2CC.tpbond.CCO2Me to N-(p-methoxyphenyl)-1-benzyl-2,3-aziridinedicarboximide was not effected thermally, but under irradsn. it gave 3 1:1-cycloadducts and a 1:2-cycloadduct I. The structural detn. of the cycloadducts was via their spectral data and chem. degradations. Mutual photochem. transformation of the cycloadducts was verified.  
 IT 25435-24-9P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of)  
 RN 25435-24-9 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-6,7-dicarboxylic acid, 8-benzyl-6-p-dioxan-2-yl-3-(p-methoxyphenyl)-2,4-dioxo-, dimethyl ester (8CI) (CA INDEX NAME)




ANSWER 33 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1969:115125 CAPLUS  
 DOCUMENT NUMBER: 70:115125  
 TITLE: Bicyclic homologs of piperazine. VIII. Synthesis and characteristics of 3,8-diazabicyclo[3.2.1]octan-2-ones  
 AUTHOR(S): Testa, Emilio; Cignarella, Giorgio; Fontanella, Luigi; Occeili, Emilio  
 CORPORATE SOURCE: Lab. Ric., "Lepetit" S.p.A., Milan, Italy  
 SOURCE: Farmaco, Edizione Scientifica (1969), 24(4), 418-34  
 CODEN: FRPSAX; ISSN: 0430-0920  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Italian  
 AB 3,8-Diazabicyclo[3.2.1]octan-2-one (I) is prepd. and converted to 3-(R-substituted)-8-(R1-substituted)-3,8-diazabicyclo[3.2.1]octan-2-ones (II). Me cis-1-carboxybenzoxy-2-carbamoylpyrrolidine-5-carboxylate (III) (m. 91-2.degree.) is prepd. from the acid. III (250 g.) is heated with 188.1 g. p-MeC6H4SO2Cl in 750 ml. pyridine to give 784 Me cis-1-carboxybenzoxy-2-cyanopyrrolidine-5-carboxylate (cis-IV)-trans-IV 1:1 mixt., b.p. 4.133-5.degree. IV (6 g.) in 100 ml. HOAc is hydrogenated over 1 g. PtO2 to give 194 I, m. 181-2.degree.; I (494) is also prepd. in the presence of Raney Ni. I (1 g.) is treated with 1 g. LiAlH4 to give 704 3,8-diazabicyclo[3.2.1]octane, b35 100.degree.; 2HCl salt m. 114-16.degree.; dipicrate m. 247-9.degree.. A mixt. of 2.6 g. I, 4 ml. 31.7% H2CO, 80 ml. water, and 2.6 g. 10% Pd/C is treated with H2 500 ml. H absorbed after 2 hrs. to give 894 II (R = CH2OH, R1 = Me) (V), b.p. 6 145.degree., citrate m. 138-40.degree.. A mixt. of 126 g. I, 126 ml. 31.5% H2CO, 250 ml. EtOH, and 126 g. 10% Pd/C is treated with H2 20 l. H is absorbed after 0.5 hr. to give 116 g. II (R = H, R1 = Me), m. 96-7.degree. (HCl salt m. 280-2.degree.), and 8 g. V. II (R = H, R1 = Me) (5 g.) is treated with 5.4 ml. 30% H2CO and 7.3 g. EtNH to give 874 3-(diethylaminomethyl)-8-methyl-3,8-diazabicyclo[3.2.1]octan-2-one, b.p. 8 144-6.degree.. Similarly prepd. are the following II (R1 = Me) (R and b.p./mm. given): 3,4,5-trimethylpiperazinomethyl, 158-60.degree./0.7; 8-propionyl-3,8-diazabicyclo[3.2.1]oct-3-ylmethyl, -. Also prepd., according to known methods, are the following II (R, R1, b.p./mm., and m.p. given): H, cyclohexylmethoxycarbonyl, -, 160-1.degree.; H, CO2CH2Ph, 100-1.degree.; COEt, COEt, -, 102-4.degree.; H, COEt, 180.degree./0.8, -, Me, COEt, 165-70.degree./0.8, 39-40.degree.; CH2CH:CHPh, COEt, 235-40.degree./0.8, -, CH2CH:CHMe2, COEt, 162-5.degree./0.6, -, Bz, Me, -, 130-2.degree.; p-ClC6H4CO, Me, -, 143-4.degree.; L-alpha-methyltropoyl, Me, -, 240-3.degree., [alpha.]20D 9.3.degree. (water); CH2CH:CHPh, Me, 194-6.degree./0.8, -, allyl, Me, 130.degree./0.8, -, p-PhC6H4CH2, Me, -, 118.degree.; propargyl, Me, 130.degree./0.8, -, H, p-PhC6H4CH2, -, 150-2.degree.; Me, p-PhC6H4CH2, -, (methiodide m. 183-4.degree.); 1.HCl, m. 262-5.degree.; 1 picrate, m. 210-12.degree..  
 IT 22315-33-9P 22315-35-1P 22315-36-2P  
 RI: SPN (Synthetic preparation); PREP (Preparation) (prepn. of)  
 RN 22315-33-9 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 8-methyl-3-(p-phenylbenzyl)- (8CI) (CA INDEX NAME)


L17 ANSWER 33 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



RN 22315-35-1 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octan-2-one, 8-(p-phenylbenzyl)- (8CI) (CA INDEX NAME)



RN 22315-36-2 CAPLUS  
 CN 3-Aza-8-azoniabicyclo[3.2.1]octane, 3,8-dimethyl-2-oxo-8-(p-phenylbenzyl)-, iodide (8CI) (CA INDEX NAME)



ANSWER 34 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1968:49651 CAPLUS  
 DOCUMENT NUMBER: 68:49651  
 TITLE: 3,8-Diastereoisomers of 3,8-diazabicyclo[3.2.1]octanes  
 INVENTOR(S): Kirchner, Frederick K.  
 PATENT ASSIGNEE(S): Sterling Drug Inc.  
 SOURCE: U.S., 8 pp.  
 CODEN: USXXAM  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 3328396		19670627	US	19611109

GI For diagram(s), see printed CA issue.  
 AB Thus, 0.1 mole diethyl 1-methyl-2,5-pyrrolidinedicarboxylate (II) was heated to 150.degree. and 0.1 mole benzylamine added during 15 min. After increasing the temp. to 180.degree., EtOH, began to distill. The temp. was increased to 280.degree. during 2 hrs. and 7.2 cc. EtOH was collected to give 3-benzyl-2,4-dioxo-3-methyl-3,8-diazabicyclo[3.2.1]octane, m. 100.0-4.4.degree. (hexane) (procedure A). The picrate m. 175-8.degree. (decompn.) (EtOH). LiAlH4 redn. of the dioxo compd. in Et2O gave I (R = PhCH2, R1 = Me) (III), b.p. 2.84-94.degree., n25D 1.5368 (procedure B). Refluxing 2.0 g. III in Et2O with excess MeI 1 hr. gave the methiodide, m. 245.0-8.2.degree. (EtOH) (procedure C). III (10 g.) in 400 cc. EtOH was acidified with concd. HCl and hydrogenated over Pd/C 6 hrs. at 23.degree. and 45 psi. to give 1.2 HCl (R = H, R1 = Me) (IV), m. 325.degree. (decompn.) (procedure D). By these procedures were prepd. 3-(3,4-dichlorobenzyl)-8-methyl-2,4-dioxo-3,8-diazabicyclo[3.2.1]octane, m. 103.6-8.6.degree. (abs. EtOH), from II and 3,4-dichlorobenzylamine; 1.2HCl (R = 3,4-Cl2C6H3CH2, R1 = Me), m. 216.4-20.8.degree. and its ethobromide 3-(4-dimethylaminobenzyl)-2,4-dioxo-8-methyl-3,8-diazabicyclo[3.2.1]octane (the di-HCl salt m. 236.8.degree. (decompn.) (dry MeOH); I (R = p-MeC6H4CH2, R1 = Me); 3-(4-chlorobenzyl)-2,4-dioxo-8-methyl-3,8-diazabicyclo[3.2.1]octane, m. 103.0-10.4.degree. (abs. EtOH); 1.2HCl (R = p-ClC6H4CH2, R1 = Me), m. 234.0-7.2.degree. and its N-benzyl chloride; 2,4-dioxo-8-methyl-3-phenethyl-3,8-diazabicyclo[3.2.1]octane HCl salt, m. 205.6-10.8.degree.; 1.2HCl (R = PhCH2CH2, R1 = Me), m. 245.degree. (decompn.) and its methosulfate; 3-(3,4-dimethylaminobenzyl)-2,4-dioxo-8-methyl-3,8-diazabicyclo[3.2.1]octane HCl salt, m. 228.degree. (decompn.); 1.2HCl (R = 3,4-(MeO)2C6H3CH2, R1 = Me), m. 194.4.degree. (decompn.) and the quaternary salt with allyl bromide. To 0.80 mole benzylamine, 0.267 mole diethyl alpha., alpha.-dibromosuccinate (V) was added with ice cooling. After warming to room temp., a vigorous reaction started and was controlled by cooling followed by heating at 95.degree.. The cryst. product was taken up in 5 N H2SO4, extd. with Et2O basified with solid NaHCO3, extd. with Et2O, the solvent evapd. in vacuo, and the residual oil distd. to give 1-benzyl-2,5-dicarbethoxypyrrolidine, b.p. 210-16.degree. (procedure E). Using procedure A, 0.05 mole of the above compd. was treated with 0.056 mole benzylamine to give 3,8-dibenzyl-2,4-dioxo-3,8-diazabicyclo[3.2.1]octane HCl salt, m. 192.4.degree.. Using procedure B, the above free base was reduced to I (R = R1 = PhCH2) (VI), m. 58.6-60.4.degree. (aq. EtOH). Using procedure C, VI was converted to the methiodide. Using excess MeI produced the bismethiodide. Using procedure D, VI was hydrogenated to give 1.2HCl (R = R1 = H), m. >300.degree. (abs. MeOH). Also, 0.027 mole IV and 0.027 mole alpha., acetoxymethylphenylacetyl chloride (VII) were added to a soln. contg. 0.08 mole NaOH in 50 cc. H2O and the mixt. kept cool overnight to


L17 ANSWER 34 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)

give 3-(3-acetoxy-2-phenylpropionyl)-8-methyl-3,8-diazabicyclo[3.2.1]octane hydrochloride (procedure F). To a soln. contg. 0.014 mole of the above salt in 50 cc. abs. MeOH, 4 cc. 4N aq. HCl was added and the soln. kept at room temp. several days. Addn. of a large amt. of abs. Et2O caused the pptn. of a white gum which solidified in abs. EtOH to give 3-(3-hydroxy-2-phenylpropionyl)-8-methyl-3,8-diazabicyclo[3.2.1]octane hydrochloride, m. 195.3-203.3.degree. (abs. EtOH-abs. Et2O). This compd. had 15.5% of the activity of atropine sulfate in the mouse after s.c. injection. The LD50 in mice was 160. +- 12 mg./kg. i.v. A soln. of 0.04 mole IV in 100 cc. dry Et2O was added dropwise over 2 hrs. to a well-stirred soln. of 0.04 mole ethanesulfonyl chloride in 100 cc. dry Et2O to give 1.HCl (R = EtSO2, R1 = Me), m. 261.2-4.6.degree. (decompn.) (MeOH). IV (0.039 mole free base) was added cautiously with cooling to 15 cc. 100% HCO2H and 0.04 mole 37% HCHO. The mixt. was heated at 95.degree. overnight, 9 cc. concd. HCl added, and the mixt. heated 3 hrs. at 95.degree. to give 1.HCl (R = R1 = Me), m. 273.2-7.4.degree. (decompn.) (EtOH). Using procedure F, 0.04 mole IV was treated with 0.04 mole VII. The acetoxy group was hydrolyzed with dil. NaOH to give 3-(4-hydroxyphenylacetyl)-8-methyl-3,8-diazabicyclo[3.2.1]octane hydrochloride, m. 226.4-7.8.degree.. Using procedure F, 0.03 mole IV was treated with 0.03 mole BzCl to give 1.HCl (R = Bz, R1 = Me), m. 264.4-7.6.degree. (abs. MeOH-Et2O). To a soln. contg. 0.012 mole of the above salt in 100 cc. abs. MeOH, 1.7 g. dry K2CO3 and 5 cc. MeI were added and the mixt. was refluxed overnight to give 1.MeI (R = Bz, R1 = Me), m. 235-8.degree.. A soln. of the methiodide in 100 cc. H2O was poured through a column of a Cl- exchange resin to give 1.MeCl (R = Bz, R1 = Me), m. 239-40.6.degree. (decompn.) (procedure G). I (R = 3-indolymethyl, R1 = Me), m. 153.8-7.60, was prepd. from 0.03 mole IV, 2.44 g. 37% HCHO, 0.03 mole indole, and 6 cc. AcOH. Excess NaOH was added to a soln. contg. 0.03 mole IV in 5 cc. H2O and the free base extd. into 200 cc. C6H6, 0.02 mole 4-chlorobenzylidene isothiocyanate in 50 cc. C6H6 added, and the soln. refluxed 1.5 hrs. to give 3-(4-chlorobenzylidenehydrazinylthiocarbonyl)-8-methyl-3,8-diazabicyclo[3.2.1]octane, m. 123.6-7.2.degree.. To a suspension of 0.12 mole IV in 60 cc. C6H6, 10 cc. 35% aq. NaOH was added. After stirring, the org. phase was sepd. and the aq. phase extd. with C6H6, the C6H6 extd. dried, and 0.02 mole diphenylchloroacetyl chloride added dropwise with stirring to give 1.HCl (R = ClCH2CO, R1 = Me), m. 248-50.degree. (decompn.) (procedure H). The above salt (5 g.) dissolved in H2O was basified with 10% Na2CO3 soln. to give I (R = HOCH2CO, R1 = Me), m. 193.6-200.6.degree.. Similarly was prepd. 1.HCl (R = Ph2CHCO, R1 = Me), m. 278.2-81.75.degree. (decompn.). To a suspension contg. 0.3 mole IV in 300 cc. C6H6, 0.7 mole NaOH and 30 cc. H2O was added. After stirring, the C6H6 phase was dried, the solvent evapd., the resultant free base dissolved in 50 cc. abs. EtOH, the soln. cooled in ice, 0.35 mole ethylene oxide in 50 cc. EtOH added, and the soln. warmed to room temp. and kept overnight to give I (R = HOCH2CH2, R1 = Me) (VIII), b.p. 89-90.degree.; di-HCl salt m. 239.4-240.8.degree. (decompn.). Using procedure A, 15 g. o-chlorobenzylamine was treated with 0.1 mole II to give 3-(2-chlorobenzyl)-8-methyl-2,4-dioxo-3,8-diazabicyclo[3.2.1]octane hydrochloride, m. 250.0-1.6.degree. (decompn.) (abs. MeOH). Using procedure B, the base corresponding to the above salt was reduced to give I (R = o-ClC6H4CH2, R1 = Me). To a soln. of 0.02 mole VIII in 20 cc. dry C6H6, 0.02 mole phenyl isocyanate was added, the soln. brought to reflux and let cool. Excess ethereal HCl was added and the resulting gum crystd. from abs. EtOH to give 3-(2-carbanilinoxyethyl)-8-methyl-3,8-diazabicyclo[3.2.1]octane as the dihydrochloride with an indefinite m.p. but softening at 176.8.degree.. To a soln. of 0.02 mole VIII and 0.02 mole NaOH in 20 cc. H2O, 0.02 mole BzCl was added dropwise, the mixt. shaken thoroughly while ice cooled, and extd. with Et2O.


L17 ANSWER 34 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
 Ethereal HCl was added to the dried Et2O ext. to give 1.2HCl [R = Et2N(CH2)2, R1 = Me], m. 197.2-209.6.degree. (decompn.). This compd. had 1.1 times the local anesthetic activity of procaine in guinea pigs. The i.v. LD50 in mice was 28 +/- 1.6 mg./kg. Using procedure E, 1.0 mole 3-diethylaminopropylamine was treated with 0.5 mole V in 700 cc. C6H6 to give 1-(3-diethylaminopropyl)-2,5-dicarbethoxypyrrolidine (IX), b0.5 135-40.degree., n25D 1.4589. Using procedure A, 0.1 mole IX was treated with 0.1 mole benzylamine to give 3-benzyl-8-(3-diethylaminopropyl)-2,4-dioxo-3,8-diazabicyclo[3.2.1]octane as the HCl salt, m. 183.0-194.8.degree. (abs. EtOH). The salt had twice the local anesthetic activity of procaine in guinea pigs. The i.v. LD50 in mice was 31 mg./kg. Using procedure B, the free base corresponding to the above salt was reduced to give I (R = PhCH2, R1 = Et2N(CH2)3). Using procedure A, 6.5 g. diethylaminopropylamine was treated with 0.05 mole II to give 3-(3-diethylaminopropyl)-8-methyl-2,4-dioxo-3,8-diazabicyclo[3.2.1]octane hydrochloride, m. 209.2-12.2.degree. (abs. EtOH). Using procedure B, the above salt was reduced to give I [R = Et2N(CH2)3, R1 = Me], b3.5 126-30.degree., n25D 1.4782; tri-HCl salt m. 169-71.degree.. Using procedure G, 0.01 mole of the above base was treated with 15 cc. MeI and 0.017 mole K2CO3 in 100 cc. abs. MeOH to give 3-(3-diethylaminopropyl)-8-methyl-3,8-diazabicyclo[3.2.1]octane bismethiodide, m. 263.4-6.4.degree. (decompn.). Using procedure A, 0.1 mole II was treated with 0.1 mole diethylaminoethylamine to give 3-(2-diethylaminoethyl)-8-methyl-2,4-dioxo-3,8-diazabicyclo[3.2.1]octane dihydrochloride, m. 221.0-2.8.degree. (decompn.). Using procedure B, the free base corresponding to the above di-HCl salt was reduced to give 1.3HCl [R = Et2NCH2CH2, R1 = Me], m. 225.6-33.2.degree. (decompn.). Using procedure A, 0.1 mole II was treated with 0.1 mole 4-diethylaminobutylamine to give 3-(4-diethylaminobutyl)-8-methyl-2,4-dioxo-3,8-diazabicyclo[3.2.1]octane dehydrochloride, m. 194.6-6.2.degree. (decompn.). Using procedure B, the above free base was reduced to give I (R = Et2N(CH2)4, R1 = Me). Using procedure A, 11.5 g. II was treated with 0.05 mole 10-(3-aminopropyl)phenothiazine to give 8-methyl-2,4-dioxo-3-[3-(10-phenothiazinyl)propyl]-3,8-diazabicyclo[3.2.1]octane, m. 133.4-7.2.degree. (95% EtOH). Using procedure B, the above base was reduced to give 1.2HCl [R = 3-(10-phenothiazinyl)propyl, R1 = Me], m. 136.2.degree. but softened at 118.0 (abs. EtOH).

IT 17740-41-9P 17740-42-0P 17783-47-0P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of)  
 RN 17740-41-9 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-2,4-dione, 3,8-bis(phenylmethyl)-, monohydrochloride (9CI) (CA INDEX NAME)


L17 ANSWER 34 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



L17 ANSWER 34 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)




● HCl  
 RN 17740-42-0 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 3,8-bis(phenylmethyl)- (9CI) (CA INDEX NAME)



RN 17783-47-0 CAPLUS  
 CN 2,5-Pyrrolidinedicarboximide, N-benzyl-1-methyl-, picrate (8CI) (CA INDEX NAME)

CM 1  
 CRN 17783-46-9  
 CMF C14 H16 N2 O2



CM 2  
 CRN 88-89-1  
 CMF C6 H3 N3 O7

L17 ANSWER 35 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN


ACCESSION NUMBER: 1964:425460 CAPLUS  
 DOCUMENT NUMBER: 61:25460  
 ORIGINAL REFERENCE NO.: 61:4374a-g  
 TITLE: Diazabicyclooctane derivatives  
 INVENTOR(S): Lepetit, S.p.A.  
 SOURCE: 12 pp.  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Unavailable  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
BE 633541		19631104	BE	
FR 1365537			FR	
GB 988526			GB	


PRIORITY APPLN. INFO.: GB 19620615

GI For diagram (p), see printed CA issue.  
 AB Improved routes to certain 3- and 8-substituted derivs. of I are described. Thus, a mixt. of 6.7 g. 3-benzyl-3,8-diazabicyclo[3.2.1]octane (II), and 12.8 g. (EtCO)2O was heated 1.5 hrs. at 100.degree., cooled, acidified with HCl, extd. with Et2O (discarded), the aq. layer alkalinized at -5.degree., and the sepd. oil extd. into Et2O. Fractionation of the ext. gave 7.4 g. 3-benzyl-8-propionyl-3,8-diazabicyclo[3.2.1]octane (III), b1 174.degree., b0.4 133-7.degree.. III was hydrogenated in 60 cc. EtOH with 3 g. 10% Pd-C at 60.degree. and 50 atm. to give 4.3 g. 8-propionyl-3,8-diazabicyclo[3.2.1]octane (IV), b0.1 121.degree.. A mixt. of 0.05 mole IV in 100 cc. Me2CO and 0.6 mole K2CO3 was stirred and treated with 0.06 mole PhCH:CHCOCl in 40 cc. Me2CO, refluxed 7 hrs., concd., the residue dissolved in 10% HCl and the soln. extd. with Et2O (discarded). The aq. layer was alkalinized, the sepd. base extd. into Et2O, and the ext. fractionated, to give 3-cinnamyl-8-propionyl-3,8-diazabicyclo[3.2.1]octane (V), b0.2 170.degree.. The following 8-propionyl derivs. were similarly prepd. (3-substituent and b.p./mm. or m.p. given): Et 85-90.degree./0.4; iso-Pr 100.degree./0.2; Bu 105.degree./0.2; allyl 95-8.degree./0.2; phenylethyl 150.degree./0.3; cyclopentylmethyl 133-8.degree./0.3; cyclopentylethyl 145.degree./0.4; phenylpropyl 175.degree./0.47; p-methoxycinnamyl 150.degree./0.1; Ph 86-8.degree.; p-nitrocinnamyl 220.degree.; p-chlorocinnamyl 220-3.degree.; o-chlorocinnamyl 201-4.degree.; p-methyldinnamyl 66-8.degree.; naphthylallyl 160-5.degree.; m-chlorocinnamyl 203-5.degree.. A soln. of 2.3 g. II in 10 cc. 2N NaOH was treated dropwise at -5.degree. with 1.92 g. EtCl, the mixt. stirred 3 hrs. at room temp., dild. with H2O, and extd. with Et2O. The ext. left 2.9 g. of viscous oil, which was treated with alc. HCl to give 3 g. HCl salt of the 3-benzyl-8-benzoyl deriv. of I, m. 219-21.degree. (EtOH). The free base, hydrogenated as described for III, gave 1.5 g. 8-benzoyl-3,8-diazabicyclo[3.2.1]octane (VI), b0.5 140-2.degree.; m. 82-3.degree. (Et2O). Similarly prepd. were the 3-benzyl-8-phenacyl (m. 78-81.degree.) and 3-benzyl-8-butyryl (m. 65-6.degree.) derivs. of I. IV was converted into its 3-propionyl isomer (VII) in 3 ways: (1) 1 g. IV was heated 5 hrs. at 120.degree. and distd. to yield 0.8 g. VII, b0.2 128-30.degree., m.p. and mixed m.p. 38-40.degree. (sublimes); (2) 1 g. IV was dissolved in 5 cc. 2N NaOH contg. a little alc., allowed to stand 4 hrs. at room temp., concd. in vacuo, and extd. with Et2O; the ext. gave 0.5 g. oil with properties as above; (3) 1 g. IV was dissolved in 10 cc. abs. alc., the soln. satd. with HCl, refluxed 2 hrs. in a current of HCl, cooled, and filtered from 0.36 g. of the 2HCl salt of I (identified by mixed m.p. and infrared spectrum). The filtrate was evapd. in vacuo, the residue alkalinized with 20% alc. NH3, Et2O added, and NH4Cl filtered off; fractionation of the filtrate gave

- L17 ANSWER 35 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
 0.58 g. oil with properties as above. To methylate VII, a soln. of 0.84 g. in 0.7 cc. HCO<sub>2</sub>H and 0.79 g. 3H<sub>2</sub>CH<sub>2</sub>O was refluxed 15 hrs., cooled, treated with 1 cc. HCl, concd. in vacuo, alkalinized, extd. with Et<sub>2</sub>O, and the ext. fractionated to give 0.54 g. 3-propionyl-8-methyl deriv. of I, b<sub>1</sub> 110-12.degree.. VII was benzylated as described for IV to V to give the 3-propionyl-8-benzyl deriv. of I, b<sub>0.2</sub> 155.degree.. VI was converted into its 3-Bz isomer [m. 122-3.degree. (Et<sub>2</sub>O)] by the 3 methods described for IV to VII.
- IT 100105-97-3, Acetophenone, 2-(3-benzyl-3,8-diazabicyclo[3.2.1]oct-8-yl)-  
 (prepn. of)
- RN 100105-97-3 CAPLUS
- CN Acetophenone, 2-(3-benzyl-3,8-diazabicyclo[3.2.1]oct-8-yl)- (7CI) (CA INDEX NAME)



- L17 ANSWER 36 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
 di-HCl salt m. 218-20.degree.; Ph, H, b<sub>0.2</sub> 114.degree.; benzyl, H, b. 95-7.degree.; di-HCl salt m. 145-8.degree.; dipicrate m. 232-5.degree.; H, H, b. 173-5.degree.; di-HCl salt m. 314-15.degree.; dipicrate m. 248-50.degree.. These compds. are pharmacologically active as diuretic, hypotensive, antihistaminic, tranquilizing, and ganglionic blocking agents.
- IT 96000-95-2, 3,8-Diazabicyclo[3.2.1]octane-8-carboxylic acid, 3-benzyl-2,4-dioxo-, benzyl ester  
 (prepn. of)
- RN 96000-95-2 CAPLUS
- CN 3,8-Diazabicyclo[3.2.1]octane-8-carboxylic acid, 3-benzyl-2,4-dioxo-, benzyl ester (6CI, 7CI) (CA INDEX NAME)



- L17 ANSWER 36 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 APPLICATION NUMBER: 1964/23448 CAPLUS  
 DOCUMENT NUMBER: 60:23448  
 ORIGINAL REFERENCE NO.: 60:4161g-h, 4162a-d  
 TITLE: 3,8-Diazabicyclo[3.2.1]octanes  
 INVENTOR(S): Cignarella, Giorgio  
 PATENT ASSIGNEE(S): Lepetit S.p.A.  
 SOURCE: 4 pp.  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Unavailable  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
GB 937184		19630918	GB	19600217


- GI For diagram(s), see printed CA Issue.
- AB The title compds. are represented by (I) wherein R is H or an alkyl (C1-8), an aryl, or an arylalkyl group, and X is H or carbobenzyloxy. The process used consists in adding to the internal anhydride of an N-substituted pyrrolidine-2,5-dicarboxylic acid an excess of an amine RNH<sub>2</sub> and refluxing the obtained crude monamine with Ac<sub>2</sub>O. Thus, a soln. of 60 g. 2,5-dicarboxy-N-benzylpyrrolidine was hydrogenated at 40.degree./20 atm. in abs. EtOH over 10% Pd-C to give 38 g. 2,5-dicarboxypyrrolidine (II), b<sub>0.3</sub> 95-6.degree.. A suspension of 200 g. II in 8 l. H<sub>2</sub>O was refluxed for 25-30 hrs. to give 110 g. pyrrolidine-2,5-dicarboxylic acid (III), m. 260-1.degree.. To a soln. of 67 g. III in 420 ml. 2N NaOH soln. cooled to 8-10.degree., 73 g. benzyl chlorocarbonate with 210 ml. 2N NaOH soln. was added with vigorous stirring during 30 min. After 2 hrs. stirring, the soln. yielded 86.5 g. N-carbobenzyloxy-pyrrolidine-2,5-dicarboxylic acid (IV), m. 125-7.degree.. A soln. of 79 g. IV in 360 ml. Ac<sub>2</sub>O was refluxed 1 hr. to give 58.1 g. IV anhydride (V), m. 166-8.degree.. To a soln. of 27.5 g. V in 300 ml. anhyd. C<sub>6</sub>H<sub>6</sub> a soln. of 1.9 g. NH<sub>3</sub> in 50 ml. C<sub>6</sub>H<sub>6</sub> was added with cooling. The mixt. was refluxed for 30 min., the solvent removed, and the resulting monamide refluxed with Ac<sub>2</sub>O 1 hr. at 130-40.degree. under 1 atm. pressure to give 18 g. 8-carbobenzyloxy-3,8-diazabicyclo[3.2.1]octane-2,4-dione (I, X = PhCH<sub>2</sub>O<sub>2</sub>C, R = H) (VI), m. 125.degree.. Similarly prepd. I (X = PhCH<sub>2</sub>O<sub>2</sub>C) analogs were (R and b.p. given): Me, b<sub>0.3</sub> 170-2.degree.; Bu, b<sub>0.3</sub> 192-4.degree.. Also prepd. were I (R, X, and m.p. given): Ph, CO<sub>2</sub>CH<sub>2</sub>Ph, 148-9.degree.; benzyl, CO<sub>2</sub>CH<sub>2</sub>Ph, 83-4.degree.; Me, H, 105-7.degree.; H, H, 223-6.degree.; benzyl, H, 78.degree. (b<sub>0.2</sub> 150-2.degree.). A mixt. of 21.5 g. II and 11.8 g. PhCH<sub>2</sub>NH<sub>2</sub> in 50 ml. dry Me<sub>2</sub>CO was refluxed for 24 hrs. to give 20.2 g. 2-carboxy-5-benzylcarbamoylpyrrolidine (VII), b<sub>0.3</sub> 178-80.degree.. Mild warming of 20.2 g. VII gave 16.5 g. 3-benzyl-3,8-diazabicyclo[3.2.1]octane-2,4-dione, b<sub>0.2</sub> 150-2.degree.. A soln. of 27.4 g. VI was added dropwise, with stirring, to an Et<sub>2</sub>O suspension of 5.7 g. LiAlH<sub>4</sub> at 0-5.degree.. The mixt. was then brought to room temp. and refluxed for 3-4 hrs. to give 8-methyl-3,8-diazabicyclo[3.2.1]octane (VIII, X = Me, R = H), b. 163-5.degree.; MeI salt m. 225-7.degree.; di-HCl salt m. 314-15.degree.; dipicrate m. 252-4.degree.. Similarly prepd. VIII were (R, X, b.p. and m.p. salts if prepd. given): Me, Me, b<sub>0.2</sub> 50-2.degree.; MeI salt m. 314-15.degree.; dipicrate m. 220-2.degree.; Ph, Me, b<sub>0.2</sub> 105-8.degree.; cinnamyl, Me, b<sub>0.6</sub> 120-30.degree.; p-chlorobenzhydryl, Me, -; HCl salt m. 176-9.degree.; o-chlorobenzhydryl, Me, -; di-HCl salt m. 283-7.degree.; benzyl, Me, -;

- L17 ANSWER 37 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN  
 APPLICATION NUMBER: 1964/23447 CAPLUS  
 DOCUMENT NUMBER: 60:23447  
 ORIGINAL REFERENCE NO.: 60:4161g-h, 4162a-d  
 TITLE: 3,8-Diazabicyclo[3.2.1]octanes  
 INVENTOR(S): Cignarella, Giorgio  
 PATENT ASSIGNEE(S): Lepetit S.p.A.  
 SOURCE: 4 pp.  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Unavailable  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
GB 937183		19630918	GB	19600217
DE 1200316			DE	
US 3221015		1965	US	

- GI For diagram(s), see printed CA Issue.
- AB The title compds. are represented by (I) wherein R is H or an alkyl (C1-8), an aryl, or an arylalkyl group, and X is H or carbobenzyloxy. The process used consists in adding to the internal anhydride of an N-substituted pyrrolidine-2,5-dicarboxylic acid an excess of an amine RNH<sub>2</sub> and refluxing the obtained crude monamine with Ac<sub>2</sub>O. Thus, a soln. of 60 g. 2,5-dicarboxy-N-benzylpyrrolidine was hydrogenated at 40.degree./20 atm. in abs. EtOH over 10% Pd-C to give 38 g. 2,5-dicarboxypyrrolidine (II), b<sub>0.3</sub> 95-6.degree.. A suspension of 200 g. II in 8 l. H<sub>2</sub>O was refluxed for 25-30 hrs. to give 110 g. pyrrolidine-2,5-dicarboxylic acid (III), m. 260-1.degree.. To a soln. of 67 g. III in 420 ml. 2N NaOH soln. cooled to 8-10.degree., 73 g. benzyl chlorocarbonate with 210 ml. 2N NaOH soln. was added with vigorous stirring during 30 min. After 2 hrs. stirring, the soln. yielded 86.5 g. N-carbobenzyloxy-pyrrolidine-2,5-dicarboxylic acid (IV), m. 125-7.degree.. A soln. of 79 g. IV in 360 ml. Ac<sub>2</sub>O was refluxed 1 hr. to give 58.1 g. IV anhydride (V), m. 166-8.degree.. To a soln. of 27.5 g. V in 300 ml. anhyd. C<sub>6</sub>H<sub>6</sub> a soln. of 1.9 g. NH<sub>3</sub> in 50 ml. C<sub>6</sub>H<sub>6</sub> was added with cooling. The mixt. was refluxed for 30 min., the solvent removed, and the resulting monamide refluxed with Ac<sub>2</sub>O 1 hr. at 130-40.degree. under 1 atm. pressure to give 18 g. 8-carbobenzyloxy-3,8-diazabicyclo[3.2.1]octane-2,4-dione (I, X = PhCH<sub>2</sub>O<sub>2</sub>C, R = H) (VI), m. 125.degree.. Similarly prepd. I (X = PhCH<sub>2</sub>O<sub>2</sub>C) analogs were (R and b.p. given): Me, b<sub>0.3</sub> 170-2.degree.; Bu, b<sub>0.3</sub> 192-4.degree.. Also prepd. were I (R, X, and m.p. given): Ph, CO<sub>2</sub>CH<sub>2</sub>Ph, 148-9.degree.; benzyl, CO<sub>2</sub>CH<sub>2</sub>Ph, 83-4.degree.; Me, H, 105-7.degree.; H, H, 223-6.degree.; benzyl, H, 78.degree. (b<sub>0.2</sub> 150-2.degree.). A mixt. of 21.5 g. II and 11.8 g. PhCH<sub>2</sub>NH<sub>2</sub> in 50 ml. dry Me<sub>2</sub>CO was refluxed for 24 hrs. to give 20.2 g. 2-carboxy-5-benzylcarbamoylpyrrolidine (VII), b<sub>0.3</sub> 178-80.degree.. Mild warming of 20.2 g. VII gave 16.5 g. 3-benzyl-3,8-diazabicyclo[3.2.1]octane-2,4-dione, b<sub>0.2</sub> 150-2.degree.. A soln. of 27.4 g. VI was added dropwise, with stirring, to an Et<sub>2</sub>O suspension of 5.7 g. LiAlH<sub>4</sub> at 0-5.degree.. The mixt. was then brought to room temp. and refluxed for 3-4 hrs. to give 8-methyl-3,8-diazabicyclo[3.2.1]octane (VIII, X = Me, R = H), b. 163-5.degree.; MeI salt m. 225-7.degree.; di-HCl salt m. 314-15.degree.; dipicrate m. 252-4.degree.. Similarly prepd. VIII were (R, X, b.p. and m.p. salts if prepd. given): Me, Me, b<sub>0.2</sub> 50-2.degree.; MeI salt m. 314-15.degree.; dipicrate m. 220-2.degree.; Ph, Me, b<sub>0.2</sub> 105-8.degree.; cinnamyl, Me, b<sub>0.6</sub> 120-30.degree.; p-chlorobenzhydryl, Me, -; HCl salt m. 176-9.degree.; o-chlorobenzhydryl, Me, -; di-HCl salt m. 283-7.degree.; benzyl, Me, -; di-HCl salt m. 218-20.degree.; Ph, H, b<sub>0.2</sub> 114.degree.; benzyl, H, b. 95-7.degree.; di-HCl salt m. 145-8.degree.; dipicrate m. 232-5.degree.; H,

L17 ANSWER 37 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
 H. b. 173-5.degree., di-HCl salt m. 314-15.degree.; dipicrate m.  
 248-50.degree.. These compds. are pharmacologically active as diuretic,  
 hypotensive, antihistaminic, tranquillizing, and ganglionic blocking  
 agents.  
 IT 96000-95-2, 3,8-Diazabicyclo[3.2.1]octane-8-carboxylic acid,  
 3-benzyl-2,4-dioxo-, benzyl ester  
 (prepn. of)  
 RN 96000-95-2 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-8-carboxylic acid, 3-benzyl-2,4-dioxo-,  
 benzyl ester (6CI, 7CI) (CA INDEX NAME)




L17 ANSWER 38 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
 233.degree.) (7 g.) in 50 cc. MeOH hydrogenated over 6 g. 10% Pd-C in 1.5  
 min. gave 93% 8-ethyl-3,8-diazabicyclo[3.2.1]octane, m. 275.degree.  
 (decompn.) (MeOH) as the di-HCl salt (XII). XII (4.3 g.) in 25 cc. Ac2O  
 and 3 g. K2CO3 heated, 12 cc. Ac2O added, refluxing continued, 5 g. K2CO3  
 added, the mixt. refluxed a few more min., cooled, MeOH added, the mixt.  
 evapd., the residue dild. with H2O, concd. KOH added, and the soln. extd.  
 with C6H6 gave 0.2 g. oil. The ext. was dissolved in 20 cc. hexane and  
 refrigerated; no solid formed, and hence the material was converted to 67%  
 HCl salt of 3-acetyl-8-ethyl-3,8-diazabicyclo[3.2.1]octane, m.  
 229-30.degree..  
 IT 17740-41-9, 2,5-Pyrrolidinedicarboximide, N,1-dibenzyl-,  
 hydrochloride 17740-42-0, 3,8-Diazabicyclo[3.2.1]octane,  
 3,8-dibenzyl- 35101-50-9, 2,5-Pyrrolidinedicarboximide,  
 1-benzoyl-N-benzyl- 102080-93-3, 3,8-Diazabicyclo[3.2.1]octane,  
 3-benzoyl-8-phenyl- 102163-86-0, 3,8-Diazabicyclo[3.2.1]octane,  
 3-benzyl-8-phenyl- 103046-68-0, 3,8-Diazabicyclo[3.2.1]octane,  
 3-benzyl-8-phenyl-, picrate 110553-68-9, 2,5-  
 Pyrrolidinedicarboximide, N-benzyl-1-phenyl- 111663-65-1,  
 3,8-Diazabicyclo[3.2.1]octane, 3,8-dibenzyl-, dihydrochloride  
 (prepn. of)  
 RN 17740-41-9 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-2,4-dione, 3,8-bis(phenylmethyl)-,  
 monohydrochloride (9CI) (CA INDEX NAME)




RN 17740-42-0 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 3,8-bis(phenylmethyl)- (9CI) (CA INDEX NAME)  
 Ph-CH2-  
 RN 35101-50-9 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane-2,4-dione, 8-benzoyl-3-(phenylmethyl)- (9CI)  
 (CA INDEX NAME)

L17 ANSWER 38 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
 ACCESSION NUMBER: 1961:144227 CAPLUS  
 DOCUMENT NUMBER: 55:144227  
 ORIGINAL REFERENCE NO.: 55:27331g-1,27332a-e  
 TITLE: Synthesis of 3,8-diazabicyclo[3.2.1]octane and some of  
 its N-substituted derivatives  
 AUTHOR(S): Blackman, Samuel W.; Baltzly, Richard  
 CORPORATE SOURCE: Polytech. Inst. of Brooklyn, Brooklyn, NY  
 SOURCE: Journal of Organic Chemistry (1961), 26, 2750-5  
 CODEN: JOCEAH; ISSN: 0022-3263  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Unavailable  
 AB The prepn. of 3,8-diazabicyclo[3.2.1]octane (I) and some of its simple  
 N-substituted derivs. was described. Adipoyl chloride brominated by a  
 standard method and added to MeOH gave (overnight) 30% di-Me  
 .alpha.,.alpha.'-dibromoadipate (II), m. 75-6.degree. (MeOH). The  
 original mother liquors and the filtrate from the crystn. of II were  
 collected and the residual ester distd. In some runs, a considerable amt.  
 of the monobromo ester was isolated as a forerun, b16, 155-70.degree..  
 The main fraction of II, b16 170-81.degree.. Recrystn. raised the yield of  
 cryst. II to 30%. II (111 g.) in 250 cc. C6H6 and 250 cc. MeNO2 refluxed  
 16 hrs. with 124 g. PhNH2 and 1 g. KI gave 27 g. crystals (two types of  
 crystals were present, long orange-yellow needles and clusters of  
 colorless crystals). Mechanical sepn. gave 13 g. trans-dimethyl  
 N-phenylpyrrolidine-2,5-dicarboxylate (III), m. 88.5.degree.  
 (Et2O-hexane); purification by chromatography on Al2O3 gave 44 g.  
 cis-ester (IV) and 13 g. III. In another run, distn. was omitted; the  
 residue dissolved in C6H6-hexane gave 67 g. mixt.; chromatography gave 14  
 g. III and 44 g. IV. IV (88 g.) in 500 cc. (CH2OH)2 was heated with 40 g.  
 PhCH2NH2 and NaOMe (72 cc. MeOH collected in the first 2 hrs.), heating  
 increased, the mixt. refluxed 16 hrs., the fractionating column removed,  
 and volatile material removed in vacuo. Distn. of the residue gave 38.7  
 g. 3-benzyl-8-phenyl-3,8-diazabicyclo[3.2.1]octane-2,4-dione (VI), m.  
 166.5.degree. (Me2CO). 8-Phenyl-3,8-diazabicyclo[3.2.1]octane-HCl (2.25  
 g.) benzoylated by the Schotten-Baumann procedure gave 1.8 g.  
 3-benzoyl-8-phenyl-3,8-diazabicyclo[3.2.1]octane (VII), m. 131-2.degree..  
 meso-II (222 g.) in 800 cc. C6H6 stirred 24 hrs. at room temp. with 225  
 cc. PhCH2NH2, the mixt. refluxed 8 hrs., cooled, and treated with Et2O,  
 the pptd. PhCH2NH2.HBr removed, the combined filtrates extd. with cold  
 dil. H2SO4, and the base liberated with Na2CO3 and extd. with Et2O gave  
 126 g. dimethyl N-benzylpyrrolidine-2,5-dicarboxylate (VIII), b1  
 148-52.degree.. VII (74 g.), 30 g. PhCH2NH2, 500 cc. (CH2OH)2, and NaOMe  
 (from 5 g. Na) gave 59 g. 3,8-dibenzyl-3,8-diazabicyclo[3.2.1]octane-2,4-  
 dione(VIII), b1 205-11.degree.. VIII (from 8.9 g. of the HCl salt) added  
 dropwise to 15.2 g. LiAlH4 in 1 l. Et2O, the mixt. stirred 30 hrs., excess  
 10% NaOH added, the mixt. stirred overnight, and the product sepd. gave  
 5.5 g. 3,8-dibenzyl-3,8-diazabicyclo[3.2.1]octane (IX), m. 52-7.degree.  
 (pentane). IX (5.15 g.) in 50 cc. MeOH contg. 5 cc. satd. MeOH-HCl  
 reduced with 6 g. 10% Pd-C and H gave 2.90 g. 1.2HCl monohydrate. I was  
 further characterized as the dipicrate. VIII (9.3 g.) in MeOH  
 hydrogenated over Pd-C gave 6.2 g. 3-benzyl-3,8-diazabicyclo[3.2.1]octane-  
 2,4-dione-HCl (X), m. 216-18.degree.. Free X (6.7 g.) benzoylated by the  
 Schotten-Baumann method and crystd. gave 3-benzyl-8-benzoyl-3,8-  
 diazabicyclo[3.2.1]octane-2,4-dione (XI), m. 121-3.degree. (with  
 frothing). XI (2 g. reduced by refluxing 24 hrs. with 2 g. LiAlH4 in Et2O  
 gave) 1 g. IX. 3-Benzyl-8-ethyl-3,8-diazabicyclo[3.2.1]octane-2HCl (m.


L17 ANSWER 38 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



RN 102080-93-3 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 3-benzoyl-8-phenyl- (6CI) (CA INDEX NAME)




RN 102163-86-0 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 3-benzyl-8-phenyl- (6CI) (CA INDEX NAME)



RN 103046-68-0 CAPLUS  
 CN 3,8-Diazabicyclo[3.2.1]octane, 3-benzyl-8-phenyl-, picrate (6CI) (CA INDEX NAME)

CH 1

CRN 102163-86-0  
 CMF C19 H22 N2




CH 2


CRN 88-89-1  
 CMF C6 H3 N3 O7




L17 ANSWER 38 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)



RN 110553-68-9 CAPLUS  
CN 2,5-Pyrrolidinedicarboximide, N-benzyl-1-phenyl- (6CI) (CA INDEX NAME)




RN 111663-65-1 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane, 3,8-dibenzyl-, dihydrochloride (6CI) (CA INDEX NAME)



● 2 HCl

L17 ANSWER 39 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN (Continued)  
ether, the solid collected, treated with a small amt. of H<sub>2</sub>O, and the undissolved product crystd. from MeOH. A soln. of 0.1 mole XI in 100 ml. anhyd. Et<sub>2</sub>O was added dropwise under stirring and cooling to a suspension of 0.5 mole LiAlH<sub>4</sub> in 200 ml. Et<sub>2</sub>O, the mixt. refluxed 4-6 hrs., cooled to 0.degree., and cautiously decompd. with 50 ml. H<sub>2</sub>O, stirred 1 hr. at room temp., filtered, the inorg. matter washed with Et<sub>2</sub>O, the ether exts. collected, dried over Na<sub>2</sub>SO<sub>4</sub> and the solvent evapd. gave I (R, 4 yield, b.p./mm., m.p. of dihydrochloride, methiodide, and dipicrate given): Me, 57, 50-2.degree./8, 260-2.degree., 250-2.degree., 242-5.degree.; Bu, 61, 54-5.degree./0.3, 245-7.degree., 218-20.degree., 220-2.degree.; Ph, 64, 104-5.degree./0.3 (m. 45-6.degree.), 180-2.degree., 262-4.degree., 172-4.degree. (monopicrate); PhCH<sub>2</sub>, 70, 88-90.degree./0.3, 213-15.degree., 250-1.degree., 228-30.degree.. A soln. of 5.4 g. I (R = CH<sub>2</sub>Ph) (XII) in 100 ml. abs. EtOH hydrogenated 2 hrs. at 50.degree. and 20 atm. in the presence of 1 g. 10% Pd-C gave 2.8 g. I (R = H) (XIII), b. 193-8.degree.; dihydrochloride m. 314-15.degree. (80% EtOH); methiodide (XIV) m. 224-5.degree.; dipicrate m. 247-50.degree.. XIV was obtained by mixing with cooling equimolar amounts of XIII (1 g.) and MeI (1.13 g.) in anhyd. Et<sub>2</sub>O and keeping 2 hrs. at room temp. The ether filtrate treated with excess MeI and kept overnight at 0.degree. gave I (R = Me) methiodide, m. 288-90.degree. (EtOH).

IT 96000-95-2, 3,8-Diazabicyclo[3.2.1]octane-8-carboxylic acid, 3-benzyl-2,4-dioxo-, benzyl ester (prepn. of)  
RN 96000-95-2 CAPLUS  
CN 3,8-Diazabicyclo[3.2.1]octane-8-carboxylic acid, 3-benzyl-2,4-dioxo-, benzyl ester (6CI, 7CI) (CA INDEX NAME)



~~L17 ANSWER 39 OF 39 CAPLUS COPYRIGHT 2003 ACS on STN~~  
~~DEPOSITION NUMBER: 1961:137533 CAPLUS~~  
~~DOCUMENT NUMBER: 55:137533~~  
~~ORIGINAL REFERENCE NO.: 55:259671,25968a-1~~

~~TITLE: Bicyclic homologs of piperazine. I. Synthesis of 8-methyl-3,8-diazabicyclooctanes~~  
~~AUTHOR(S): Cignarella, Giorgio; Nathansohn, Giangiacomo~~  
~~CORPORATE SOURCE: Lepetit S.p.A., Milan~~  
~~SOURCE: Journal of Organic Chemistry (1961), 26, 1500-4~~  
~~CODEN: JOCEAH; ISSN: 0022-3263~~

~~DOCUMENT TYPE: Journal~~  
~~LANGUAGE: Unavailable~~  
~~GI For diagram(s), see printed CA issue.~~

~~AB The synthesis and properties of 3-substituted 8-methyl-3,8-diazabicyclo[3.2.1]octanes (I), in which the 3-substituent was H, Me, Bu, Ph, and CH<sub>2</sub>Ph, were described. To a refluxed soln. of 500 g. di-Et meso-.alpha.,.alpha.-dibromoadipate (II) in 1500 ml. C<sub>6</sub>H<sub>6</sub>, 490 g. PhCH<sub>2</sub>NH<sub>2</sub> (III) was added under stirring in 1 hr., the mixt. refluxed 24 hrs. while PhCH<sub>2</sub>NH<sub>2</sub>.HBr sepd., cooled, the ppt. filtered off, washed with C<sub>6</sub>H<sub>6</sub>, the C<sub>6</sub>H<sub>6</sub> soln. evapd. in vacuo, and the oily residue distd. to give 350 g. cis-2,5-dicarbethoxy-N-benzylpyrrolidine (IV), b<sub>0.3</sub> 145-8.degree., HCl salt m. 123-5.degree. (EtOH). A soln. of 60 g. IV in 600 ml. abs. EtOH was hydrogenated at 20 atm./40.degree. in the presence of 9 g. 10% Pd-C 2 hrs. to give 38 g. cis-2,5-dicarbethoxypyrrolidine (V), b<sub>0.3</sub> 95-6.degree., HCl salt m. 134-5.degree. (EtOH-Et<sub>2</sub>O). A suspension of 200 g. V in 8 l. H<sub>2</sub>O refluxed under stirring 25-30 hrs. gave 98 g. cis-pyrrolidine-2,5-dicarboxylic acid (VI), m. 260-1.degree.. VI (67 g.) added to a 420 ml. 2N NaOH at -10.degree. was treated under stirring and cooling with 75 g. PhCH<sub>2</sub>COCl (VII) 30 min. and the mixt. stirred 2 hrs. at room temp. to yield 94 g. cis-N-carbobenzoxypyrrolidine-2,5-dicarboxylic acid (VIII), m. 125-7.degree. (H<sub>2</sub>O). VIII (79 g.) and 360 ml. Ac<sub>2</sub>O refluxed 1 hr. the solvent removed in vacuo and the residue kept at 140.degree./1 mm. 1 hr., 100 ml. anhyd. AcOEt added to the solid residue, the mixt. kept overnight at room temp., and the solid collected, washed with cold AcOEt, and dried yielded 58.1 g. (crude) N-carbobenzoxypyrrolidine-2,5-dicarboxylic acid anhydride (IX), m. 170-1.degree. (C<sub>6</sub>H<sub>6</sub> or Me<sub>2</sub>CO). To 0.1 mole IX in 300 ml. anhyd. C<sub>6</sub>H<sub>6</sub>, 0.11 mole of appropriate amine in 50 ml. C<sub>6</sub>H<sub>6</sub> was added with cooling, the mixt. heated on a water bath 20 min., and the solvent evapd. to give the crude 2N.CH(CONHR).CH<sub>2</sub>.CH<sub>2</sub>.CHCO<sub>2</sub>H (X) (Z = carbobenzoxo throughout; R = Me, Bu, Ph, CH<sub>2</sub>Ph) (Xa, Xb, Xc, and Xd, resp.). X (R = H), m. 179-81.degree., was obtained in 80% yield (a) by satg. with NH<sub>3</sub> a C<sub>6</sub>H<sub>6</sub> soln. of IX, evapg. the C<sub>6</sub>H<sub>6</sub>, dissolving the residue in H<sub>2</sub>O, and cautiously acidifying with 10% HCl; the oil sepd. solidified and crystd. from H<sub>2</sub>O to give 23.4 g. of the desired product; (b) by adding IX to 33% NH<sub>4</sub>OH to 0.degree., evapg. excess ammonia, and acidifying at pH 4 with 10% HCl. Xa-Xb refluxed with 5 vols. Ac<sub>2</sub>O 1-2 hrs., the solvent distd. and the residual oil kept 1-2 hrs. at 130-40.degree./1 mm., cooled, 200 ml. 1:1 Et<sub>2</sub>O:C<sub>6</sub>H<sub>6</sub> added, the insol. oil sepd., the clear soln. washed with 5% NaHCO<sub>3</sub> and H<sub>2</sub>O, dried over Na<sub>2</sub>SO<sub>4</sub>, and the solvent evapd. gave the following CH<sub>2</sub>.CH<sub>2</sub>.CH.NZ.CH.CO.NR.CO (XI) (R, 4 yield from IX, b.p./mm. or m.p. and solvent of crystn. given): H, 50.4, 124-5.degree., MeOH; Me, 60, 170-2.degree./0.3; Bu, 63, 192-4.degree./0.3; Ph, 45, 148-9.degree., MeOH; CH<sub>2</sub>Ph, 64, 83-4.degree., MeOH. XI (R = Ph) was also obtained in 39.7% yield by adding a suspension of 20 g. IX in 60 ml. pyridine to BzN<sub>3</sub> in 60 ml. pyridine; the mixt. was heated at 100-10.degree. 1 hr. till gas evolution subsided, the pyridine evapd. in vacuo, the residue taken up in~~